

The Guide to Clinical Preventive Services

2014



AHRQ
Agency for Healthcare Research and Quality
Advancing Excellence in Health Care • www.ahrq.gov

 U.S. Preventive Services
TASK FORCE

The clinical summaries in the Guide are abridged versions of recommendations from the U.S. Preventive Services Task Force (USPSTF). To view the full recommendation statements, supporting evidence, or recommendations published after March 2014, go to www.USPreventiveServicesTaskForce.org.

The USPSTF Electronic Preventive Services Selector (ePSS) allows users to download the USPSTF recommendations to PDA, mobile, or tablet devices; receive notifications of updates; and search and browse recommendations online. Users can search the ePSS for recommendations by patient age, sex, and pregnancy status. To download, subscribe, or search, go to epss.ahrq.gov.

Recommendations made by the USPSTF are independent of the U.S. Government. They should not be construed as an official position of AHRQ or the U.S. Department of Health and Human Services.

The Guide to Clinical Preventive Services 2014

Recommendations of the U.S.
Preventive Services Task Force



Agency for Healthcare Research and Quality
Advancing Excellence in Health Care • www.ahrq.gov

Foreword

Since 1998, the Agency for Healthcare Research and Quality (AHRQ) has convened the U.S. Preventive Services Task Force (USPSTF)—an independent, volunteer panel of national experts in prevention and evidence-based medicine. The Task Force makes recommendations to help primary care clinicians and patients decide together whether a preventive service is right for a patient’s needs.

AHRQ staff provide scientific and administrative support for the Task Force, and assist in disseminating its findings and recommendations to key audiences. In that role, we are pleased to make *The Guide to Clinical Preventive Services 2014* available to those who seek to ensure that their patients receive the highest quality clinical preventive services.

I am gratified that AHRQ has been able to support the USPSTF in its efforts to engage the public and to ensure that its processes are transparent. Most notably, all Task Force draft materials are now available for public comment online at www.USPreventiveServicesTaskForce.org.

Previous iterations of the USPSTF *Guide to Clinical Preventive Services* are used around the Nation to provide appropriate and effective preventive care. The *Guide* is designed to be user-friendly for practicing clinicians. In addition, the *Guide* lists resources that clinicians can use to educate their patients on appropriate preventive services, such as *Stay Healthy* brochures and MyHealthfinder (see Appendixes). Also included are brief descriptions of and links to tools that health care professionals can use to improve their practice, including the electronic Preventive Services Selector (ePSS) and the National Guideline Clearinghouse (see Appendixes).

As more information becomes available to clinicians and patients alike, AHRQ’s goal is to help improve patients’ health and well-being, and contribute to better health outcomes for the Nation overall.

Richard Kronick, Ph.D.
Director
Agency for Healthcare Research and Quality

Preface

The U.S. Preventive Services Task Force (USPSTF) is mandated by Congress to conduct rigorous reviews of scientific evidence to create evidence-based recommendations for preventive services that may be provided in the primary care setting.

Since its inception, the USPSTF has made and maintained recommendations on dozens of clinical preventive services that are intended to prevent or reduce the risk for heart disease, cancer, infectious diseases, and other conditions and events that impact the health of children, adolescents, adults, and pregnant women. The *Guide to Clinical Preventive Services 2014* includes both new and updated recommendations released from 2004-2014 in a brief, easily usable format meant for use at the point of patient care. The most up-to-date version of the recommendations, as well as the complete USPSTF recommendation statements, are available along with their supporting scientific evidence at www.USPreventiveServicesTaskForce.org.

Recommendations for preventive care have evolved over time as it has become widely recognized that some “preventive” services were not actually beneficial. Individual health care providers, professional organizations, integrated health systems, insurers (both private and public), as well as groups crafting health quality measures and national health objectives, have recognized the need to carefully balance potential benefits and harms using the highest quality of evidence, and have adopted the recommendations of the Task Force. The primary audience for the USPSTF’s work remains primary care clinicians, and the recommendations are now considered by many to provide definitive standards for preventive services. The work of the USPSTF is recognized by the Patient Protection and Affordable Care Act. Under the law, preventive services with a Task Force grade of A or B must be covered without cost-sharing (e.g., copayment or deductible) under new health insurance plans or policies.

In the last several years, the USPSTF has increased the transparency of its work, and these efforts have gained additional momentum in view of the enhanced importance of the recommendations under the new law. Public comments are welcomed at multiple points in the development of each recommendation to encourage additional input from experts, advocates and other stakeholders to help the Task Force craft relevant and clear recommendation statements. However, the USPSTF remains committed to evaluating evidence free from the influence of politics, special interests, and advocacy.

As the science around clinical practice guideline development has evolved, USPSTF methods continue to evolve as well. The Procedure Manual, which can be found on

the USPSTF Web site, details the most updated version of the process for evaluating the quality and strength of the evidence for a service, determining the net health benefit (benefit minus harms) associated with the service, and judging the level of certainty that providing these services will be beneficial in primary care. We continue to explore the appropriate use of mathematical modeling to help fill research gaps regarding the ages at which to start and stop providing a service, and at what time intervals. In addition, we are committed to improving the communication of our recommendations to a broader audience, including patients and policymakers.

The letter grade linked to each recommendation reflects both the magnitude of net benefit and the strength and certainty of the evidence supporting the provision of a specific preventive service. These grades translate to practice guidance for clinicians:

- Discuss services with “A” and “B” recommendation grades with eligible patients and offer them as a priority.
- Discourage the use of services with “D” recommendation grades unless there are unusual additional considerations.
- Give lower priority to services with “C” recommendation grades; they need not be provided unless there are individual considerations in favor of providing the service.
- Help patients understand the uncertainty surrounding services with “I” (insufficient evidence) statements, which reflect the conclusion that the evidence is insufficient to determine net benefit. The Clinical Considerations section of each full recommendation statement offers additional guidance.

The USPSTF recognizes that clinical decisions about patients involve more complex considerations than the evidence alone; clinicians should always understand the evidence but individualize decisionmaking to the specific patient and situation. The Clinical Considerations section of each USPSTF recommendation statement helps clinicians by offering practical information so they can tailor these recommendations to individual patients.

We strongly encourage clinicians to visit the USPSTF Web site and read the complete recommendation statements for services relevant to their patients. Additional information is available to facilitate the delivery of the highest quality preventive care. For each topic, educational materials have been developed for use with patients and the public. Special materials developed for clinicians are also available on some complex topics, such as prostate and lung cancer screening, along with links to informative Web sites. In addition, the USPSTF Electronic Preventive Services

Selector (ePSS), available via PDA, smart phone, or on the Web at epss.ahrq.gov, allows users to search USPSTF recommendations by patient age and other clinical characteristics.

We hope you find the *Guide to Clinical Preventive Services 2014* to be a useful tool as you care for patients. Based on the best medical evidence available, we are confident that by implementing these recommended services, you will help your patients live longer and healthier lives.

Michael L. LeFevre, M.D., M.S.P.H., Chair
Albert L. Siu, M.D., M.S.P.H., Co-Vice Chair
Kirsten Bibbins-Domingo, Ph.D., M.D., M.A.S., Co-Vice Chair
U.S. Preventive Services Task Force

Contents

| | |
|---|-----|
| Foreword | iii |
| Preface | v |
| Preventive Services Recommended by the USPSTF | 1 |
| Clinical Summaries of Recommendations for Adults (alphabetical list) | 5 |
| Abdominal Aortic Aneurysm, Screening..... | 7 |
| *Alcohol Misuse, Screening and Behavioral Counseling | 8 |
| Aspirin for the Prevention of Cardiovascular Disease, Preventive Medication | 9 |
| Aspirin or NSAIDS for Prevention of Colorectal Cancer, Preventive Medication.... | 10 |
| Bacterial Vaginosis in Pregnancy, Screening | 11 |
| Bacteriuria, Screening..... | 12 |
| Bladder Cancer, Screening..... | 13 |
| *BRCA-Related Cancer in Women, Screening | 14 |
| *Breast Cancer, Preventive Medications..... | 15 |
| Breast Cancer, Screening | 16 |
| Breastfeeding, Counseling | 17 |
| Carotid Artery Stenosis, Screening | 18 |
| Cervical Cancer, Screening..... | 19 |
| Chlamydial Infection, Screening | 20 |
| *Chronic Kidney Disease, Screening | 21 |
| Chronic Obstructive Pulmonary Disease, Screening..... | 22 |
| *Cognitive Impairment in Older Adults, Screening..... | 23 |
| Colorectal Cancer, Screening..... | 24 |
| Coronary Heart Disease (Risk Assessment, Nontraditional Risk Factors), Screening..... | 25 |
| *Coronary Heart Disease, Screening (With Electrocardiography)..... | 26 |
| Depression in Adults, Screening | 27 |
| Diabetes Mellitus, Screening | 28 |
| *Falls in Older Adults, Counseling, Preventive Medication, and Other Interventions | 29 |
| Folic Acid Supplementation to Prevent Neural Tube Defects, Preventive Medication | 30 |
| Genital Herpes, Screening..... | 31 |
| *Gestational Diabetes Mellitus, Screening | 32 |
| *Glaucoma, Screening | 33 |
| Gonorrhea, Screening..... | 34 |
| *Healthful Diet and Physical Activity, Counseling..... | 35 |
| *Hearing Loss in Older Adults, Screening | 36 |

| | |
|---|----|
| Hemochromatosis, Screening | 37 |
| Hepatitis B Virus Infection in Pregnant Women, Screening | 38 |
| *Hepatitis C Virus Infection in Adults, Screening | 9 |
| High Blood Pressure in Adults, Screening..... | 40 |
| *HIV Infection, Screening..... | 41 |
| Illicit Drug Use, Screening | 42 |
| Impaired Visual Acuity in Older Adults, Screening | 43 |
| *Intimate Partner Violence and Elderly Abuse, Screening..... | 44 |
| Lipid Disorders in Adults, Screening | 45 |
| *Lung Cancer, Screening | 46 |
| *Menopausal Hormone Therapy for the Primary Prevention of Chronic Conditions, Preventive Medication | 47 |
| Motor Vehicle Occupant Restraints, Counseling | 48 |
| *Obesity in Adults, Screening and Counseling | 49 |
| *Oral Cancer, Screening | 50 |
| Osteoporosis, Screening..... | 51 |
| *Ovarian Cancer, Screening..... | 52 |
| *Peripheral Artery Disease and Cardiovascular Risk Assessment, Screening | 53 |
| *Prostate Cancer, Screening..... | 54 |
| Sexually Transmitted Infections, Counseling | 55 |
| *Skin Cancer, Counseling..... | 56 |
| Skin Cancer, Screening..... | 57 |
| Suicide Risk, Screening..... | 58 |
| Syphilis Infection (Pregnant Women), Screening..... | 59 |
| Testicular Cancer, Screening..... | 60 |
| Tobacco Use in Adults, Counseling and Interventions..... | 61 |
| *Vitamin D and Calcium Supplementation to Prevent Fractures, Preventive Medication | 62 |
| *Vitamin, Mineral, and Multivitamin Supplements for Primary Prevention of Cardiovascular Disease and Cancer, Preventive Medication | 63 |

| | |
|---|-----------|
| Clinical Summaries of Recommendations for Children and Adolescents (alphabetical list) | 65 |
| *Alcohol Misuse, Screening and Behavioral Counseling | 8 |
| Blood Lead Levels in Children and Pregnant Women, Screening | 67 |
| Cervical Cancer, Screening | 19 |
| *Child Maltreatment, Interventions to Prevent | 68 |
| Chlamydial Infection, Screening | 20 |
| Congenital Hypothyroidism, Screening..... | 69 |
| Developmental Dysplasia of the Hip, Screening..... | 70 |
| Genital Herpes, Screening | 31 |
| Gonococcal Ophthalmia Neonatorum, Preventive Medication..... | 71 |

| | |
|--|-----------|
| Gonorrhea, Screening..... | 34 |
| Hearing Loss in Newborns, Screening..... | 72 |
| *High Blood Pressure in Children, Screening..... | 73 |
| *HIV Infection, Screening..... | 41 |
| Hyperbilirubinemia in Infants, Screening..... | 74 |
| *Illicit and Prescription Drug Use in Children and Adolescents, Counseling..... | 75 |
| Iron Deficiency Anemia, Screening..... | 76, 77 |
| Lipid Disorders in Children, Screening..... | 78 |
| Major Depressive Disorder in Children and Adolescents, Screening..... | 79 |
| Motor Vehicle Occupant Restraints, Counseling..... | 48 |
| Obesity in Children and Adolescents, Screening..... | 80 |
| Phenylketonuria (PKU), Screening..... | 81 |
| Sexually Transmitted Infections, Counseling..... | 55 |
| Sickle Cell Disease in Newborns, Screening..... | 82 |
| *Skin Cancer, Counseling..... | 56 |
| Speech and Language Delay, Screening..... | 83 |
| Suicide Risk, Screening..... | 58 |
| Testicular Cancer, Screening..... | 60 |
| *Tobacco Use in Children and Adolescents, Primary Care Interventions..... | 84 |
| Visual Impairment in Children Ages 1-5, Screening..... | 85 |
| Immunizations..... | 87 |
| Topics in Progress..... | 91 |
| Appendixes and Index..... | 95 |
| Appendix A. How the U.S. Preventive Services Task Force Grades Its Recommendations..... | 97 |
| Appendix B. Members of the U.S. Preventive Services Task Force 2004-2014..... | 101 |
| Appendix C. Acknowledgements..... | 104 |
| Appendix D. About the U.S. Preventive Services Task Force..... | 107 |
| Appendix E. More Resources..... | 112 |
| Appendix F. 2002 Breast Cancer Screening Recommendation..... | 115 |
| Index. Recommendations, Cross-Referenced..... | 117 |

*New recommendations released March 2012 to March 2014.



Preventive Services Recommended by the USPSTF

All clinical summaries in this Guide are abridged recommendations. To see the full recommendation statements and recommendations published after March 2014, go to www.USPreventiveServicesTaskForce.org.

Section 1: Preventive Services Recommended by the USPSTF

The U.S. Preventive Services Task Force (USPSTF) recommends that clinicians discuss these preventive services with eligible patients and offer them as a priority. All these services have received an “A” or a “B” (recommended) grade from the Task Force. Refer to the endnotes for each recommendation for population-specific clinical considerations.

For definitions of all grades used by the USPSTF, see Appendix A (beginning on p. 97). The full listings of all USPSTF recommendations for adults begin on p. 5 and recommendations for children begin on p. 65.

| Recommendation | Adults | | | Special Populations | | |
|---|---|-------|--|---------------------|-----------------------|---|
| | Men | Women | | Pregnant Women | Children/ Adolescents | |
| | Abdominal Aortic Aneurysm, Screening ¹ | ✓ | | | | |
| Alcohol Misuse Screening and Behavioral Counseling | ✓ | ✓ | | ✓ | | |
| Aspirin for the Prevention of Cardiovascular Disease ² | ✓ | ✓ | | | | |
| Bacteriuria, Screening ³ | | | | ✓ | | |
| BRCA-Related Cancer in Women, Screening ⁴ | | ✓ | | | | |
| Breast Cancer, Preventive Medications ⁵ | | ✓ | | | | |
| Breast Cancer, Screening ⁶ | | ✓ | | | | |
| Breastfeeding, Counseling ⁷ | | ✓ | | ✓ | | |
| Cervical Cancer, Screening ⁸ | | ✓ | | | | |
| Chlamydial Infection, Screening ⁹ | | ✓ | | ✓ | | |
| Colorectal Cancer, Screening ¹⁰ | ✓ | ✓ | | | | |
| Congenital Hypothyroidism, Screening ¹¹ | | | | | | ✓ |
| Depression in Adults, Screening ¹² | ✓ | ✓ | | | | |
| Diabetes Mellitus, Screening ¹³ | ✓ | ✓ | | | | |
| Falls in Older Adults, Counseling, Preventive Medication, and Other Interventions ¹⁴ | ✓ | ✓ | | | | |

| Section 1: Preventive Services Recommended by the USPSTF (continued) | | | | | | |
|--|--------|-------|---------------------|---|--------------------------|---|
| Recommendation | Adults | | Special Populations | | Children/ Adolescents | |
| | Men | Women | Pregnant Women | | | |
| Folic Acid Supplementation to Prevent Neural Tube Defects, Preventive Medication ¹⁵ | | ✓ | | | | |
| Gestational Diabetes Mellitus, Screening ¹⁶ | | | ✓ | | | |
| Gonococcal Ophthalmia Neonatorum, Preventive Medication ¹⁷ | | | | | ✓ | |
| Gonorrhea, Screening ¹⁸ | | ✓ | | | | |
| Hearing Loss in Newborns, Screening ¹⁹ | | | | | | ✓ |
| Hepatitis B Virus Infection in Pregnant Women, Screening ²⁰ | | | ✓ | | | |
| Hepatitis C Virus Infection in Adults, Screening ²¹ | ✓ | | ✓ | | | ✓ |
| High Blood Pressure in Adults, Screening | ✓ | | | | | |
| HIV Infection, Screening ²² | ✓ | | ✓ | | | ✓ |
| Intimate Partner Violence and Elderly Abuse, Screening ²³ | | | ✓ | | | |
| Iron Deficiency Anemia, Prevention ²⁴ | | | | | | ✓ |
| Iron Deficiency Anemia, Screening ²⁵ | | | | ✓ | | |
| Lipid Disorders in Adults, Screening ²⁶ | ✓ | | ✓ | | | |
| Lung Cancer, Screening ²⁷ | ✓ | | ✓ | | | |
| Major Depressive Disorder in Children and Adolescents, Screening ²⁸ | | | | | | ✓ |
| Obesity in Adults, Screening ²⁹ | ✓ | | ✓ | | | |
| Obesity in Children and Adolescents, Screening ³⁰ | | | | | | ✓ |
| Osteoporosis, Screening ³¹ | | | ✓ | | | |

| Section 1: Preventive Services Recommended by the USPSTF (continued) | | | | | |
|---|--------|-------|---------------------|----------------------|---|
| Recommendation | Adults | | Special Populations | | |
| | Men | Women | Pregnant Women | Children/Adolescents | |
| Phenylketonuria (PKU), Screening ³² | | | | ✓ | |
| Sexually Transmitted Infections, Counseling ³³ | ✓ | ✓ | | ✓ | |
| Sickle Cell Disease in Newborns, Screening ³⁴ | | | | ✓ | |
| Skin Cancer, Counseling ³⁵ | ✓ | ✓ | ✓ | ✓ | |
| Syphilis Infection (Pregnant Women), Screening | | | ✓ | | |
| Tobacco Use in Adults, Counseling and Interventions ³⁶ | ✓ | ✓ | ✓ | | |
| Tobacco Use in Children and Adolescents, Primary Care Interventions ³⁷ | | | | ✓ | |
| Visual Impairment in Children Ages 1 to 5, Screening ³⁸ | | | | | ✓ |

¹One-time screening by ultrasonography in men aged 65 to 75 who have ever smoked.

²When the potential harm of an increase in gastrointestinal hemorrhage is outweighed by a potential benefit of a reduction in myocardial infarctions (men aged 45-79 years) or in ischemic strokes (women aged 55-79 years).

³Pregnant women at 12-16 weeks gestation or at first prenatal visit, if later.

⁴Refer women whose family history is associated with an increased risk for deleterious mutations in *BRCA1* or *BRCA2* genes for genetic counseling and evaluation for *BRCA* testing.

⁵Engage in shared, informed decisionmaking and offer to prescribe risk-reducing medications, if appropriate, to women aged ≥35 years without prior breast cancer diagnosis who are at increased risk.

⁶Biennial screening mammography for women aged 50 to 74 years. Note: The Department of Health and Human Services, in implementing the Affordable Care Act, follows the 2002 USPSTF recommendation for screening

mammography, with or without clinical breast examination, every 1-2 years for women aged 40 and older.

⁷Interventions during pregnancy and after birth to promote and support breastfeeding.

⁸Screen with cytology every 3 years (women ages 21 to 65) or co-test (cytology/HPV testing) every 5 years (women ages 30-65).

⁹Sexually active women 24 and younger and other asymptomatic women at increased risk for infection. Asymptomatic pregnant women 24 and younger and others at increased risk.

¹⁰Adults aged 50-75 using fecal occult blood testing, sigmoidoscopy, or colonoscopy.

¹¹Newborns.

¹²When staff-assisted depression care supports are in place to assure accurate diagnosis, effective treatment, and followup.

- ¹³Asymptomatic adults with sustained blood pressure greater than 135/80 mg Hg.
- ¹⁴Provide intervention (exercise or physical therapy and/or vitamin D supplementation) to community-dwelling adults ≥65 years at increased risk for falls.
- ¹⁵All women planning or capable of pregnancy take a daily supplement containing 0.4 to 0.8 mg (400 to 800 µg) of folic acid.
- ¹⁶Asymptomatic pregnant women after 24 weeks of gestation.
- ¹⁷Newborns.
- ¹⁸Sexually active women, including pregnant women 25 and younger, or at increased risk for infection.
- ¹⁹Newborns.
- ²⁰Screen at first prenatal visit.
- ²¹Persons at high risk for infection and adults born between 1945 and 1965.
- ²²All adolescents and adults ages 15 to 65 years and others who are at increased risk for HIV infection and all pregnant women.
- ²³Asymptomatic women of childbearing age; provide or refer women who screen positive to intervention services.
- ²⁴Routine iron supplementation for asymptomatic children aged 6 to 12 months who are at increased risk for iron deficiency anemia.
- ²⁵Routine screening in asymptomatic pregnant women.
- ²⁶Men aged 20-35 and women over age 20 who are at increased risk for coronary heart disease; all men aged 35 and older.
- ²⁷Asymptomatic adults aged 55 to 80 years who have a 30 pack-year smoking history and currently smoke or have quit smoking within the past 15 years.
- ²⁸Adolescents (age 12 to 18) when systems are in place to ensure accurate diagnosis, psychotherapy, and followup.
- ²⁹Patients with a body mass index of 30 kg/m² or higher should be offered or referred to intensive, multicomponent behavioral interventions.
- ³⁰Screen children aged 6 years and older; offer or refer for intensive counseling and behavioral interventions.
- ³¹Women aged 65 years and older and women under age 65 whose 10-year fracture risk is equal to or greater than that of a 65-year-old white woman without additional risk factors.
- ³²Newborns.
- ³³All sexually active adolescents and adults at increased risk for STIs.
- ³⁴Newborns.
- ³⁵Children, adolescents, and young adults aged 10 to 24 years.
- ³⁶Ask all adults about tobacco use and provide tobacco cessation interventions for those who use tobacco; provide augmented, pregnancy-tailored counseling for those pregnant women who smoke.
- ³⁷Provide interventions to prevent initiation of tobacco use in school-aged children and adolescents.
- ³⁸Screen children ages 3 to 5 years.



Clinical Summaries of Recommendations for Adults

All clinical summaries in this Guide are abridged recommendations. To see the full recommendation statements and recommendations published after March 2014, go to www.USPreventiveServicesTaskForce.org.

Abdominal Aortic Aneurysm

| Screening for Abdominal Aortic Aneurysm | |
|--|---|
| Population | Men ages 65 to 75 years who have ever smoked Women ages 65 to 75 years |
| Recommendation | Screen once for abdominal aortic aneurysm with ultrasonography. Grade: B No recommendation for or against screening. Grade: C Do not screen for abdominal aortic aneurysm. Grade: D |
| Risk Assessment | The major risk factors for abdominal aortic aneurysm include male sex, a history of ever smoking (defined as 100 cigarettes in a person's lifetime), and age of 65 years or older. |
| Screening Tests | Screening abdominal ultrasonography is an accurate test when performed in a setting with adequate quality assurance (i.e., in an accredited facility with credentialed technologists). Abdominal palpation has poor accuracy and is not an adequate screening test. |
| Timing of Screening | One-time screening to detect an abdominal aortic aneurysm using ultrasonography is sufficient. There is negligible health benefit in re-screening those who have normal aortic diameter on initial screening. |
| Interventions | Open surgical repair of an aneurysm of at least 5.5 cm leads to decreased abdominal aortic aneurysm-related mortality in the long term; however, there are major harms associated with this procedure. |
| Balance of Benefits and Harms | In men ages 65 to 75 years who have ever smoked, the benefits of screening for abdominal aortic aneurysm outweigh the harms. In men ages 65 to 75 years who have never smoked, the balance between the benefits and harms of screening for abdominal aortic aneurysm is too close to make a general recommendation for this population. The potential overall benefit of screening for abdominal aortic aneurysm among women ages 65 to 75 years is low because of the small number of abdominal aortic aneurysm-related deaths in this population and the harms associated with surgical repair. |
| Other Relevant USPSTF Recommendations | The USPSTF has made recommendations on screening for carotid artery stenosis, coronary heart disease, high blood pressure, lipid disorders, and peripheral arterial disease. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Alcohol Misuse

| Screening and Behavioral Counseling Interventions in Primary Care To Reduce Alcohol Misuse | |
|---|---|
| Title | Adolescents |
| Population | Adults aged 18 years or older |
| Recommendation | <p>Screen for alcohol misuse and provide brief behavioral counseling interventions to persons engaged in risky or hazardous drinking.</p> <p>Grade: B</p> <p>No recommendation.</p> <p>Grade: I (Insufficient Evidence)</p> |

| | |
|--|---|
| Screening Tests | <p>Numerous screening instruments can detect alcohol misuse in adults with acceptable sensitivity and specificity. The USPSTF prefers the following tools for alcohol misuse screening in the primary care setting:</p> <ol style="list-style-type: none"> 1. AUDIT 2. Abbreviated AUDIT-C 3. Single-question screening, such as asking, "How many times in the past year have you had 5 (for men) or 4 (for women and all adults older than 65 years) or more drinks in a day?" |
| Behavioral Counseling Interventions | <p>Counseling interventions in the primary care setting can improve unhealthy alcohol consumption behaviors in adults engaging in risky or hazardous drinking. Behavioral counseling interventions for alcohol misuse vary in their specific components, administration, length, and number of interactions. Brief multicontact behavioral counseling seems to have the best evidence of effectiveness; very brief behavioral counseling has limited effect.</p> |
| Balance of Benefits and Harms | <p>There is a moderate net benefit to alcohol misuse screening and brief behavioral counseling interventions in the primary care setting for adults aged 18 years or older.</p> <p>The evidence on alcohol misuse screening and brief behavioral counseling interventions in the primary care setting for adolescents is insufficient, and the balance of benefits and harms cannot be determined.</p> |
| Other Relevant USPSTF Recommendations | <p>The USPSTF has made recommendations on screening for illicit drug use and counseling and interventions to prevent tobacco use. These recommendations are available at http://www.uspreventiveservicestaskforce.org/.</p> |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Aspirin for the Prevention of Cardiovascular Disease

| Aspirin for the Prevention of Cardiovascular Disease | | | | | |
|--|---|---|---|---|----------------------------------|
| Title | Aspirin for the Prevention of Cardiovascular Disease | | | | |
| Population | Men age 45-79 years | Women age 55-79 years | Men age <45 years | Women age <55 years | Men Women age ≥80 years |
| Recommendation | Encourage aspirin use when potential CVD benefit (MIs prevented) outweighs potential harm of GI hemorrhage. | Encourage aspirin use when potential CVD benefit (strokes prevented) outweighs potential harm of GI hemorrhage. | Do not encourage aspirin use for MI prevention. | Do not encourage aspirin use for stroke prevention. | No Recommendation |
| | Grade: A | | Grade: D | | Grade: I (Insufficient Evidence) |

| <p>How to Use This Recommendation</p> <p>Shared decisionmaking is strongly encouraged with individuals whose risk is close to (either above or below) the estimates of 10-year risk levels indicated below. As the potential CVD benefit increases above harms, the recommendation to take aspirin should become stronger.</p> <p>To determine whether the potential benefit of MIs prevented (men) and strokes prevented (women) outweighs the potential harm of increased GI hemorrhage, both 10-year CVD risk and age must be considered.</p> <p>Risk level at which CVD events prevented (benefit) exceeds GI harms</p> | <table border="1" style="width: 100%; text-align: center;"> <thead> <tr> <th colspan="2">Men</th> <th colspan="2">Women</th> </tr> <tr> <th colspan="2">10-year CHD risk</th> <th colspan="2">10-year stroke risk</th> </tr> </thead> <tbody> <tr> <td>Age 45-59 years</td> <td>≥4%</td> <td>Age 55-59 years</td> <td>≥3%</td> </tr> <tr> <td>Age 60-69 years</td> <td>≥9%</td> <td>Age 60-69 years</td> <td>≥8%</td> </tr> <tr> <td>Age 70-79 years</td> <td>≥12%</td> <td>Age 70-79 years</td> <td>≥11%</td> </tr> </tbody> </table> | | | Men | | Women | | 10-year CHD risk | | 10-year stroke risk | | Age 45-59 years | ≥4% | Age 55-59 years | ≥3% | Age 60-69 years | ≥9% | Age 60-69 years | ≥8% | Age 70-79 years | ≥12% | Age 70-79 years | ≥11% |
|---|--|-----------------|---------------------|-----|--|-------|--|------------------|--|---------------------|--|-----------------|-----|-----------------|-----|-----------------|-----|-----------------|-----|-----------------|------|-----------------|------|
| | Men | | Women | | | | | | | | | | | | | | | | | | | | |
| | 10-year CHD risk | | 10-year stroke risk | | | | | | | | | | | | | | | | | | | | |
| | Age 45-59 years | ≥4% | Age 55-59 years | ≥3% | | | | | | | | | | | | | | | | | | | |
| Age 60-69 years | ≥9% | Age 60-69 years | ≥8% | | | | | | | | | | | | | | | | | | | | |
| Age 70-79 years | ≥12% | Age 70-79 years | ≥11% | | | | | | | | | | | | | | | | | | | | |
| <p>The table above applies to adults who are not taking NSAIDs and who do not have upper GI pain or a history of GI ulcers.</p> <p>NSAID use and history of GI ulcers raise the risk of serious GI bleeding considerably and should be considered in determining the balance of benefits and harms. NSAID use combined with aspirin use approximately quadruples the risk of serious GI bleeding compared to the risk with aspirin use alone. The rate of serious bleeding in aspirin users is approximately 2-3 times higher in patients with a history of GI ulcers.</p> | | | | | | | | | | | | | | | | | | | | | | | |
| <p>For men: Risk factors for CHD include age, diabetes, total cholesterol level, HDL level, blood pressure, and smoking. CHD risk estimation tool: cvdrisk.nhlbi.nih.gov/calculator.asp</p> <p>For women: Risk factors for ischemic stroke include age, high blood pressure, diabetes, smoking, history of CVD, atrial fibrillation, and left ventricular hypertrophy. Stroke risk estimation tool: http://www.westernstroke.org/index.php?header_name_stroke_tools.gif_main_stroke_tools.php</p> | | | | | | | | | | | | | | | | | | | | | | | |
| <p>Risk Assessment</p> <p>The USPSTF has made recommendations on screening for abdominal aortic aneurysm, carotid artery stenosis, coronary heart disease, high blood pressure, lipid disorders, and peripheral arterial disease. These recommendations are available at http://www.uspreventiveservicestaskforce.org.</p> | | | | | | | | | | | | | | | | | | | | | | | |

| |
|--|
| <p>Other Relevant USPSTF Recommendations</p> <p>The USPSTF has made recommendations on screening for abdominal aortic aneurysm, carotid artery stenosis, coronary heart disease, high blood pressure, lipid disorders, and peripheral arterial disease. These recommendations are available at http://www.uspreventiveservicestaskforce.org.</p> |
|--|

Abbreviations: CHD coronary heart disease, CVD cardiovascular disease, GI gastrointestinal, HDL high-density lipoprotein, MI myocardial infarction, NSAIDs nonsteroidal anti-inflammatory drugs.

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Aspirin or NSAIDs for Prevention Of Colorectal Cancer

| | |
|--|---|
| Title | Routine Aspirin or Nonsteroidal Anti-Inflammatory Drug (NSAID) Use for the Primary Prevention of Colorectal Cancer |
| Population | Asymptomatic adults at average risk for colorectal cancer |
| Recommendation | Do not use aspirin or NSAIDs for the prevention of colorectal cancer. Grade: D |
| Risk Assessment | The major risk factors for colorectal cancer are older age (older than age 50 years), family history (having two or more first- or second-degree relatives with colorectal cancer), and African American race. |
| Balance of Benefits and Harms | Aspirin and NSAIDs, taken in higher doses for longer periods, reduce the incidence of adenomatous polyps. However, there is poor evidence that aspirin and NSAID use leads to a reduction in colorectal cancer-associated mortality. Aspirin increases the incidence of gastrointestinal bleeding and hemorrhagic stroke; NSAIDs increase the incidence of gastrointestinal bleeding and renal impairment, especially in the elderly. The USPSTF concluded that the harms outweigh the benefits of aspirin and NSAID use for the prevention of colorectal cancer. |
| Other Relevant USPSTF Recommendations | The USPSTF has made recommendations on screening for colorectal cancer and aspirin use for the prevention of cardiovascular disease. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Bacterial Vaginosis in Pregnancy

| Title | Screening for Bacterial Vaginosis in Pregnancy to Prevent Preterm Delivery | |
|-----------------------|--|--|
| Population | Asymptomatic pregnant women without risk factors for preterm delivery | Asymptomatic pregnant women with risk factors for preterm delivery |
| Recommendation | Do not screen. Grade: D | No recommendation. Grade: I (Insufficient Evidence) |

| | |
|----------------------------|--|
| Risk Assessment | <p>Risk factors of preterm delivery include:</p> <ul style="list-style-type: none"> ● African-American women. ● Pelvic infection. ● Previous preterm delivery. <p>Bacterial vaginosis is more common among African-American women, women of low socioeconomic status, and women who have previously delivered low-birth-weight infants.</p> <p>Bacterial vaginosis is diagnosed using Amsel's clinical criteria or Gram stain.</p> <p>When using Amsel's criteria, 3 out of 4 criteria must be met to make a clinical diagnosis:</p> <ol style="list-style-type: none"> 1. Vaginal pH >4.7. 2. The presence of clue cells on wet mount. 3. Thin homogeneous discharge. 4. Amine 'fishy odor' when potassium hydroxide is added to the discharge. |
| Screening Tests | <p>Not applicable.</p> |
| Screening Intervals | <p>Treatment is appropriate for pregnant women with symptomatic bacterial vaginosis infection.</p> <p>Oral metronidazole and oral clindamycin, as well as vaginal metronidazole gel or clindamycin cream, are used to treat bacterial vaginosis.</p> <p>The optimal treatment regimen is unclear.¹</p> |

¹The Centers for Disease Control and Prevention (CDC) recommends 250 mg oral metronidazole 3 times a day for 7 days as the treatment for bacterial vaginosis in pregnancy.

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Bacteriuria

| Screening for Asymptomatic Bacteriuria in Adults | |
|---|--|
| Title | |
| Population | All pregnant women Men and nonpregnant women |
| Recommendation | Screen with urine culture. Grade: A Do not screen. Grade: D |
| Detection and Screening Tests | Asymptomatic bacteriuria can be reliably detected through urine culture. The presence of at least 10 ⁵ colony-forming units per mL of urine, of a single uropathogen, and in a midstream clean-catch specimen is considered a positive test result. |
| Screening Intervals | A clean-catch urine specimen should be collected for screening culture at 12-16 weeks' gestation or at the first prenatal visit, if later. The optimal frequency of subsequent urine testing during pregnancy is uncertain. |
| Benefits of Detection and Early Treatment | The detection and treatment of asymptomatic bacteriuria with antibiotics significantly reduces the incidence of symptomatic maternal urinary tract infections and low birthweight. |
| Harms of Detection and Early Treatment | Potential harms associated with treatment of asymptomatic bacteriuria include: <ul style="list-style-type: none"> ● Adverse effects from antibiotics. ● Development of bacterial resistance. |
| Other Relevant USPSTF Recommendations | Additional USPSTF recommendations involving screening for infectious conditions during pregnancy can be found at www.uspreventiveservicestaskforce.org/recommendations.htm#obstetric and www.uspreventiveservicestaskforce.org/recommendations.htm#infectious . |

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Bladder Cancer

| | |
|--|--|
| Title | Screening for Bladder Cancer |
| Population | Asymptomatic adults |
| Recommendation | No recommendation. Grade: I (Insufficient Evidence) |
| Risk Assessment | <p>Risk factors for bladder cancer include:</p> <ul style="list-style-type: none"> ● Smoking ● Occupational exposure to carcinogens (e.g., rubber, chemical, and leather industries) ● Male sex ● Older age ● White race ● Infections caused by certain bladder parasites ● Family or personal history of bladder cancer |
| Screening Tests | <p>Screening tests for bladder cancer include:</p> <ul style="list-style-type: none"> ● Microscopic urinalysis for hematuria ● Urine cytology ● Urine biomarkers |
| Interventions | <p>The principal treatment for superficial bladder cancer is transurethral resection of the bladder tumor, which may be combined with adjuvant radiation therapy, chemotherapy, biologic therapies, or photodynamic therapies. Radical cystectomy, often with adjuvant chemotherapy, is used in cases of surgically resectable invasive bladder cancer.</p> |
| Balance of Benefits and Harms | <p>There is inadequate evidence that treatment of screen-detected bladder cancer leads to improved morbidity or mortality. There is inadequate evidence on harms of screening for bladder cancer.</p> |
| Suggestions for Practice | <p>In deciding whether to screen for bladder cancer, clinicians should consider the following:</p> <ul style="list-style-type: none"> ● <i>Potential preventable burden</i>: early detection of tumors with malignant potential could have an important impact on the mortality rate of bladder cancer. ● <i>Potential harms</i>: false-positive results may lead to anxiety and unneeded evaluations, diagnostic-related harms from cystoscopy and biopsy, harms from labeling and unnecessary treatments, and overdiagnosis. ● <i>Current practice</i>: screening tests used in primary practice include microscopic urinalysis for hematuria and urine cytology; urine biomarkers are not commonly used in part because of cost. Patients with positive findings are typically referred to a urologist for further evaluation. |
| Other Relevant USPSTF Recommendations | <p>Recommendations on screening for other types of cancer can be found at www.uspreventiveservicestaskforce.org.</p> |

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

BRCA-Related Cancer in Women

| | | |
|--|---|---|
| Title | Risk Assessment, Genetic Counseling, and Genetic Testing for BRCA-Related Cancer in Women | |
| Population | Asymptomatic women who have not been diagnosed with BRCA-related cancer | |
| Recommendation | <p>Screen women whose family history may be associated with an increased risk for potentially harmful BRCA mutations. Women with positive screening results should receive genetic counseling and, if indicated after counseling, BRCA testing.</p> <p>Grade: B</p> | <p>Do not routinely recommend genetic counseling or BRCA testing to women whose family history is not associated with an increased risk for potentially harmful BRCA mutations.</p> <p>Grade: D</p> |
| Risk Assessment | <p>Family history factors associated with increased likelihood of potentially harmful BRCA mutations include breast cancer diagnosis before age 50 years, bilateral breast cancer, family history of breast and ovarian cancer, presence of breast cancer in ≥ 1 male family member, multiple cases of breast cancer in the family, ≥ 1 or more family member with 2 primary types of BRCA-related cancer, and Ashkenazi Jewish ethnicity. Several familial risk stratification tools are available to determine the need for in-depth genetic counseling, such as the Ontario Family History Assessment Tool, Manchester Scoring System, Referral Screening Tool, Pedigree Assessment Tool, and FHS-7.</p> <p>Genetic risk assessment and BRCA mutation testing are generally multistep processes involving identification of women who may be at increased risk for potentially harmful mutations, followed by genetic counseling by suitably trained health care providers and genetic testing of selected high-risk women when indicated.</p> <p>Tests for BRCA mutations are highly sensitive and specific for known mutations, but interpretation of results is complex and generally requires posttest counseling.</p> | |
| Screening Tests | <p>Interventions in women who are BRCA mutation carriers include earlier, more frequent, or intensive cancer screening; risk-reducing medications (e.g., tamoxifen or raloxifene); and risk-reducing surgery (e.g., mastectomy or salpingo-oophorectomy).</p> | |
| Treatment | <p>In women whose family history is associated with an increased risk for potentially harmful BRCA mutations, the net benefit of genetic testing and early intervention is moderate.</p> <p>In women whose family history is not associated with an increased risk for potentially harmful BRCA mutations, the net benefit of genetic testing and early intervention ranges from minimal to potentially harmful.</p> | |
| Balance of Benefits and Harms | <p>The USPSTF has made recommendations on medications for the reduction of breast cancer risk and screening for ovarian cancer. These recommendations are available at www.uspreventiveservicestaskforce.org.</p> | |
| Other Relevant USPSTF Recommendations | <p>The USPSTF has made recommendations on medications for the reduction of breast cancer risk and screening for ovarian cancer. These recommendations are available at www.uspreventiveservicestaskforce.org.</p> | |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to www.uspreventiveservicestaskforce.org.

Breast Cancer (Preventive Medications)

| Title | Medications for Risk Reduction of Primary Breast Cancer in Women | |
|--|--|--|
| Population | Asymptomatic women aged ≥35 years without a prior diagnosis of breast cancer who are at increased risk for the disease | Asymptomatic women aged ≥35 years without a prior diagnosis of breast cancer who are not at increased risk for the disease |
| Recommendation | Engage in shared, informed decision making and offer to prescribe risk-reducing medications, if appropriate. Grade: B | |
| Risk Assessment | <p>Important risk factors for breast cancer include patient age, race/ethnicity, age at menarche, age at first live childbirth, personal history of ductal or lobular carcinoma in situ, number of first-degree relatives with breast cancer, personal history of breast biopsy, body mass index, menopausal status or age, breast density, estrogen and progestin use, smoking, alcohol use, physical activity, and diet.</p> <p>Available risk assessment models can accurately predict the number of breast cancer cases that may arise in certain study populations, but their ability to accurately predict which women will develop breast cancer is modest.</p> | |
| Preventive Medications | <p>The selective estrogen receptor modulators tamoxifen and raloxifene have been shown to reduce the incidence of invasive breast cancer in women who are at increased risk for the disease. Tamoxifen has been approved for this use in women age 35 years or older, and raloxifene has been approved for this use in postmenopausal women. The usual daily doses for tamoxifen and raloxifene are 20 mg and 60 mg, respectively, for 5 years.</p> | |
| Balance of Benefits and Harms | There is a moderate net benefit from use of tamoxifen and raloxifene to reduce the incidence of invasive breast cancer in women who are at increased risk for the disease. | The potential harms of tamoxifen and raloxifene outweigh the potential benefits for breast cancer risk reduction in women who are not at increased risk for the disease. Potential harms include thromboembolic events, endometrial cancer, and cataracts. |
| Other Relevant USPSTF Recommendations | The USPSTF has made recommendations on risk assessment, genetic counseling, and genetic testing for BRCA-related cancer, as well as screening for breast cancer. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . | |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Breast Cancer (Screening)

| Screening for Breast Cancer: Using Film Mammography | | | |
|--|---|--|---|
| Title | Women aged 40-49 years | Women aged 50-74 years | Women aged ≥75 years |
| Population | Women aged 40-49 years | Women aged 50-74 years | Women aged ≥75 years |
| Recommendation | Individualize decision to begin biennial screening according to the patient's circumstances and values. Grade: C | Screen every 2 years. Grade: B | No recommendation. Grade: I (Insufficient Evidence) |
| Risk Assessment | This recommendation applies to women aged ≥40 years who are not at increased risk by virtue of a known genetic mutation or history of chest radiation. Increasing age is the most important risk factor for most women. | | |
| Screening Tests | Standardization of film mammography has led to improved quality. Refer patients to facilities certified under the Mammography Quality Standards Act (MQSA), listed at http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfm/QSA/mqsa.cfm | | |
| Timing of Screening | Evidence indicates that biennial screening is optimal. A biennial schedule preserves most of the benefit of annual screening and cuts the harms nearly in half. A longer interval may reduce the benefit. | | |
| Balance of Benefits and Harms | <p>There is convincing evidence that screening with film mammography reduces breast cancer mortality, with a greater absolute reduction for women aged 50 to 74 years than for younger women.</p> <p>Harms of screening include psychological harms, additional medical visits, imaging, and biopsies in women without cancer; inconvenience due to false-positive screening results; harms of unnecessary treatment, and radiation exposure. Harms seem moderate for each age group.</p> <p>False-positive results are a greater concern for younger women; treatment of cancer that would not become clinically apparent during a woman's life (overdiagnosis) is an increasing problem as women age.</p> | | |
| Rationale for No Recommendation (I Statement) | Among women 75 years or older, evidence of benefit is lacking. | | |
| Other Relevant USPSTF Recommendations | USPSTF recommendations on screening for genetic susceptibility for breast cancer and chemoprevention of breast cancer are available at http://www.uspreventiveservicestaskforce.org . | | |

The U.S. Department of Health and Human Services, in implementing the Affordable Care Act under the standard it sets out in revised Section 2713(a)(5) of the Public Health Service Act, utilizes the 2002 recommendation on breast cancer screening of the U.S. Preventive Services Task Force. For clinical summary of 2002 Recommendation, see Appendix F.

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Breastfeeding

| Primary Care Interventions to Promote Breastfeeding | | | |
|---|---|---|--|
| Title | Primary Care Interventions to Promote Breastfeeding | | |
| Population | Pregnant women | New mothers | The mother's partner, other family members, and friends |
| Recommendation | <p>Promote and support breastfeeding.</p> <p>Grade: B</p> | | |
| Benefits of Breastfeeding | <p>Mothers</p> <p>Less likelihood of breast and ovarian cancer</p> | <p>Infants</p> <p>Fewer ear infections, lower-respiratory-tract infections, and gastrointestinal infections</p> | <p>Young children</p> <p>Less likelihood of asthma, type 2 diabetes, and obesity</p> |
| Interventions to Promote Breastfeeding | <p>Interventions to promote and support breastfeeding have been found to increase the rates of initiation, duration, and exclusivity of breastfeeding. Consider multiple strategies, including:</p> <ul style="list-style-type: none"> ● Formal breastfeeding education for mothers and families ● Direct support of mothers during breastfeeding ● Training of primary care staff about breastfeeding and techniques for breastfeeding support ● Peer support <p>Interventions that include both prenatal and postnatal components may be most effective at increasing breastfeeding duration. In rare circumstances, for example for mothers with HIV and infants with galactosemia, breastfeeding is not recommended. Interventions to promote breastfeeding should empower individuals to make informed choices supported by the best available evidence.</p> | | |
| Implementation | System-level interventions with senior leadership support may be more likely to be sustained over time. | | |

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Carotid Artery Stenosis

| | |
|---|---|
| Title | Screening for Carotid Artery Stenosis |
| Population | Adult general population ¹ |
| Recommendation | Do not screen with ultrasound or other screening tests. Grade: D |
| Risk Assessment | The major risk factors for carotid artery stenosis (CAS) include: older age, male gender, hypertension, smoking, hypercholesterolemia, and heart disease. However, accurate, reliable risk assessment tools are not available. |
| Balance of Benefits and Harms | Harms outweigh benefits. In the general population, screening with carotid duplex ultrasound would result in more false-positive results than true positive results. This would lead either to surgeries that are not indicated or to confirmatory angiography. As the result of these procedures, some people would suffer serious harms (death, stroke, and myocardial infarction) that outweigh the potential benefit surgical treatment may have in preventing stroke. |
| Other Relevant Recommendations from the USPSTF | Adults should be screened for hypertension, hyperlipidemia, and smoking. Clinicians should discuss aspirin chemoprevention with patients at increased risk for cardiovascular disease. These recommendations and related evidence are available at http://www.uspreventiveservicestaskforce.org . |

¹This recommendation applies to adults without neurological symptoms and without a history of transient ischemic attacks (TIA) or stroke. If otherwise eligible, an individual who has a carotid area TIA should be evaluated promptly for consideration of carotid endarterectomy.

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Cervical Cancer

| Screening for Cervical Cancer | |
|-------------------------------|--|
| Title | Screening for Cervical Cancer |
| Population | Women ages 21 to 65 |
| Recommendation | Screen with cytology (Pap smear) every 3 years. Grade: A |
| | Women ages 30 to 65 |
| | Screen with cytology every 3 years or co-testing (cytology/HPV testing) every 5 years Grade: A |
| | Women younger than age 21 |
| | Do not screen. Grade: D |
| | Women older than age 65 who have had adequate prior screening and are not high risk |
| | Do not screen. Grade: D |
| | Women after hysterectomy with removal of the cervix and with no history of high-grade precancer or cervical cancer |
| | Do not screen. Grade: D |
| | Women younger than age 30 |
| | Do not screen with HPV testing (alone or with cytology) Grade: D |

| | | | | | | |
|---|--|---|--|---|---|---|
| Risk Assessment | Human papillomavirus (HPV) infection is associated with nearly all cases of cervical cancer. Other factors that put a woman at increased risk of cervical cancer include HIV infection, a compromised immune system, in utero exposure to diethylstilbestrol, and previous treatment of a high-grade precancerous lesion or cervical cancer. | | | | | |
| Screening Tests and Interval | Screening women ages 21 to 65 years every 3 years with cytology provides a reasonable balance between benefits and harms. Screening with cytology more often than every 3 years confers little additional benefit, with large increases in harms. HPV testing combined with cytology (co-testing) every 5 years in women ages 30 to 65 years offers a comparable balance of benefits and harms, and is therefore a reasonable alternative for women in this age group who would prefer to extend the screening interval. | | | | | |
| Timing of Screening | Screening earlier than age 21 years, regardless of sexual history, leads to more harms than benefits. Clinicians and patients should base the decision to end screening on whether the patient meets the criteria for adequate prior testing and appropriate follow-up, per established guidelines. | | | | | |
| Interventions | Screening aims to identify high-grade precancerous cervical lesions to prevent development of cervical cancer and early-stage asymptomatic invasive cervical cancer. High-grade lesions may be treated with ablative and excisional therapies, including cryotherapy, laser ablation, loop excision, and cold knife conization. Early-stage cervical cancer may be treated with surgery (hysterectomy) or chemotherapy. | | | | | |
| Balance of Benefits and Harms | <table border="1" style="width: 100%;"> <tbody> <tr> <td>The benefits of screening with cytology every 3 years substantially outweigh the harms.</td> <td>The benefits of screening with co-testing (cytology/HPV testing) every 5 years outweigh the harms.</td> <td>The harms of screening earlier than age 21 years outweigh the benefits.</td> <td>The benefits of screening after age 65 years do not outweigh the potential harms.</td> <td>The potential harms of screening with HPV testing (alone or with cytology) outweigh the potential benefits.</td> </tr> </tbody> </table> | The benefits of screening with cytology every 3 years substantially outweigh the harms. | The benefits of screening with co-testing (cytology/HPV testing) every 5 years outweigh the harms. | The harms of screening earlier than age 21 years outweigh the benefits. | The benefits of screening after age 65 years do not outweigh the potential harms. | The potential harms of screening with HPV testing (alone or with cytology) outweigh the potential benefits. |
| The benefits of screening with cytology every 3 years substantially outweigh the harms. | The benefits of screening with co-testing (cytology/HPV testing) every 5 years outweigh the harms. | The harms of screening earlier than age 21 years outweigh the benefits. | The benefits of screening after age 65 years do not outweigh the potential harms. | The potential harms of screening with HPV testing (alone or with cytology) outweigh the potential benefits. | | |
| Other Relevant USPSTF Recommendations | The USPSTF has made recommendations on screening for breast cancer and ovarian cancer, as well as genetic risk assessment and <i>BRCA</i> mutation testing for breast and ovarian cancer susceptibility. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . | | | | | |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Chlamydial Infection

| Title | Screening for Chlamydial Infection | | | | | |
|----------------|--|--|----------------------|--|---------------------|--|
| | Non-pregnant women | | Pregnant women | | Men | |
| Population | 24 years and younger | 25 years and older | 24 years and younger | 25 years and older | Men | |
| | Includes adolescents | Not at increased risk | Includes adolescents | Not at increased risk | At increased risk | |
| Recommendation | Screen if sexually active. Grade: A | Do not automatically screen. Grade: C | Screen. Grade: A | Do not automatically screen. Grade: C | Screen. Grade: B | No recommendation. Grade: I (Insufficient Evidence ¹) |

| | |
|----------------------------|---|
| Risk Assessment | <p>Age: Women and men aged 24 years and younger are at greatest risk.</p> <p>History of: previous chlamydial infection or other sexually transmitted infections, new or multiple sexual partners, inconsistent condom use, sex work.</p> <p>Demographics: African-Americans and Hispanic women and men have higher prevalence rates than the general population in many communities.</p> |
| Screening Tests | Nucleic acid amplification tests (NAATs) can identify chlamydial infection in asymptomatic women (non-pregnant and pregnant) and asymptomatic men. NAATs have high specificity and sensitivity and can be used with urine and vaginal swabs. |
| Screening Intervals | <p>Non-Pregnant Women The optimal interval for screening is not known. The CDC recommends that women at increased risk be screened at least annually.²</p> <p>Pregnant Women For women 24 years and younger and older women at increased risk: Screen at the first prenatal visit. For patients at continuing risk, or who are newly at risk: Screen in the 3rd trimester.</p> |
| Treatment | The Centers for Disease Control and Prevention has outlined appropriate treatment at: http://www.cdc.gov/STD/treatment . Test and/or treat partners of patients treated for chlamydial infection. |

¹Chlamydial infection results in few sequelae in men. Therefore, the major benefit of screening men would be to reduce the likelihood that infected and untreated men would pass the infection to sexual partners. There is no evidence that screening men reduces the long-term consequences of chlamydial infection in women. Because of this lack of evidence, the USPSTF was not able to assess the balance of benefits and harms, and concluded that the evidence is insufficient to recommend for or against routinely screening men.

²Centers for Disease Control and Prevention. Sexually transmitted diseases treatment guidelines, 2006. *MMWR* 2006; 55(No. RR-11).

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Chronic Kidney Disease

| | |
|--|---|
| Title | Screening for Chronic Kidney Disease |
| Population | Asymptomatic adults without diagnosed chronic kidney disease (CKD) |
| Recommendation | No recommendation. Grade: I (Insufficient Evidence) |
| Risk Assessment | There is no generally accepted risk assessment tool for CKD or risk for complications of CKD. Diabetes and hypertension are well-established risk factors with strong links to CKD. Other risk factors for CKD include older age, cardiovascular disease, obesity, and family history. |
| Screening Tests | While there is insufficient evidence to recommend routine screening, the tests often suggested for screening that are feasible in primary care include testing the urine for protein (microalbuminuria or macroalbuminuria) and testing the blood for serum creatinine to estimate glomerular filtration rate. |
| Balance of Benefits and Harms | The USPSTF could not determine the balance between the benefits and harms of screening for CKD in asymptomatic adults. |
| Other Relevant USPSTF Recommendations | The USPSTF has made recommendations on screening for diabetes, hypertension, and obesity, as well as aspirin use for the prevention of cardiovascular disease. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Chronic Obstructive Pulmonary Disease

| | |
|--|---|
| Title | Screening for Chronic Obstructive Pulmonary Disease Using Spirometry |
| Population | Adult general population |
| Recommendation | Do not screen for chronic obstructive pulmonary disease using spirometry. Grade: D |
| Additional Population Information | This screening recommendation applies to healthy adults who do not recognize or report respiratory symptoms to a clinician. It does not apply to individuals with a family history of alpha-1 antitrypsin deficiency. |
| Risk Assessment | Risk factors for COPD include: <ul style="list-style-type: none"> ● Current or past tobacco use. ● Exposure to occupational and environmental pollutants. ● Age 40 or older. |
| Screening Tests¹ | Spirometry can be performed in a primary care physician's office or a pulmonary testing laboratory. The USPSTF did not review evidence comparing the accuracy of spirometry performed in primary care versus referral settings. For individuals who present to clinicians complaining of chronic cough, increased sputum production, wheezing, or dyspnea, spirometry would be indicated as a diagnostic test for COPD, asthma, and other pulmonary diseases. |
| Other Approaches to the Prevention of Pulmonary Illnesses | These services should be offered to patients regardless of COPD status: <ul style="list-style-type: none"> ● All current smokers should receive smoking cessation counseling and be offered pharmacologic therapies demonstrated to increase cessation rates. ● All patients 50 years of age or older should be offered influenza immunization annually. ● All patients 65 years of age or older should be offered one-time pneumococcal immunization. |
| Other Relevant USPSTF Recommendations | Clinicians should screen all adults for tobacco use and provide tobacco cessation interventions for those who use tobacco products. The USPSTF tobacco cessation counseling recommendation and supporting evidence are available at http://www.uspreventiveservicestaskforce.org/uspstf/uspstbac.htm . |

¹The potential benefit of spirometry-based screening for COPD is prevention of one or more exacerbations by treating patients found to have an airflow obstruction previously undetected. However, even in groups with the greatest prevalence of airflow obstruction, hundreds of patients would need to be screened with spirometry to defer one exacerbation.

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Cognitive Impairment

| Screening for Cognitive Impairment in Older Adults | |
|---|--|
| Title | |
| Population | Community-dwelling adults who are older than age 65 years and have no signs or symptoms of cognitive impairment |
| Recommendation | No recommendation. Grade: I statement |
| Risk Assessment | Increasing age is the strongest known risk factor for cognitive impairment. Other reported risk factors for cognitive impairment include cardiovascular risk factors (such as diabetes, tobacco use, hypercholesterolemia, and hypertension), head trauma, learning disabilities (such as Down syndrome), depression, alcohol abuse, physical frailty, low education level, low social support, and having never been married. |
| Screening Tests | Screening tests for cognitive impairment in the clinical setting generally include asking patients to perform a series of tasks that assess 1 or more cognitive domains (memory, attention, language, and visuospatial or executive functioning). The most widely studied instrument is the Mini-Mental State Examination. Other instruments with more limited evidence include the Clock Draw Test, Mini-Cog, Memory Impairment Screen, Abbreviated Mental Test, Short Portable Mental Status Questionnaire, Free and Cued Selective Reminding Test, 7-Minute Screen, Telephone Interview for Cognitive Status, and Informant Questionnaire on Cognitive Decline in the Elderly. |
| Treatment | Pharmacologic treatments approved by the U.S. Food and Drug Administration include acetylcholinesterase inhibitors and memantine. Nonpharmacologic interventions include cognitive training, lifestyle behavioral interventions, exercise, educational interventions, and multidisciplinary care interventions. Some interventions focus on the caregiver and aim to improve caregiver morbidity and delay institutionalization of persons with dementia. |
| Balance of Benefits and Harms | The evidence on screening for cognitive impairment is lacking, and the balance of benefits and harms cannot be determined. |
| Other Relevant USPSTF Recommendations | The USPSTF has made recommendations related to several of the risk factors for cognitive impairment, including counseling on tobacco cessation, alcohol use, healthful diet, physical activity, and falls prevention and screening for high cholesterol, hypertension, and depression. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Colorectal Cancer

| Screening for Colorectal Cancer | | | |
|---|--|--|----------------------------|
| Title | | | |
| Population¹ | Adults age 50 to 75 years | Adults age 76 to 85 years | Adults older than 85 |
| Recommendation | Screen with high sensitivity fecal occult blood testing (FOBT), sigmoidoscopy, or colonoscopy. Grade: A | Do not automatically screen. Grade: C | Do not screen. Grade: D |
| <p>For all populations, evidence is insufficient to assess the benefits and harms of screening with computerized tomography colonography (CTC) and fecal DNA testing. Grade: I (Insufficient Evidence)</p> | | | |

| | | | |
|--|--|---|--|
| Screening Tests | <p>High sensitivity FOBT, sigmoidoscopy with FOBT, and colonoscopy are effective in decreasing colorectal cancer mortality. The risks and benefits of these screening methods vary. Colonoscopy and flexible sigmoidoscopy (to a lesser degree) entail possible serious complications.</p> | | |
| Screening Test Intervals | <p>Intervals for recommended screening strategies:</p> <ul style="list-style-type: none"> ● Annual screening with high-sensitivity fecal occult blood testing ● Sigmoidoscopy every 5 years, with high-sensitivity fecal occult blood testing every 3 years ● Screening colonoscopy every 10 years | | |
| Balance of Benefits and Harms | The benefits of screening outweigh the potential harms for 50- to 75-year-olds. | <p>The likelihood that detection and early intervention will yield a mortality benefit declines after age 75 because of the long average time between adenoma development and cancer diagnosis.</p> | |
| Implementation | <p>Focus on strategies that maximize the number of individuals who get screened. Practice shared decisionmaking; discussions with patients should incorporate information on test quality and availability. Individuals with a personal history of cancer or adenomatous polyps are followed by a surveillance regimen, and screening guidelines are not applicable.</p> | | |
| Other Relevant USPSTF Recommendations | <p>The USPSTF recommends against the use of aspirin or nonsteroidal anti-inflammatory drugs for the primary prevention of colorectal cancer. This recommendation is available at http://www.uspreventiveservicestaskforce.org.</p> | | |

¹These recommendations do not apply to individuals with specific inherited syndromes (Lynch Syndrome or Familial Adenomatous Polyposis) or those with inflammatory bowel disease.

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Coronary Heart Disease (Risk Assessment, Nontraditional Risk Factors)

| | |
|--|---|
| Title | Using Nontraditional Risk Factors In Coronary Heart Disease Risk Assessment |
| Population | Asymptomatic men and women with no history of coronary heart disease (CHD), diabetes, or any CHD risk equivalent |
| Recommendation | No recommendation. Grade: I (Insufficient Evidence) |
| Risk Assessment | This recommendation applies to adult men and women classified at intermediate 10-year risk for CHD (10% to 20%) by traditional risk factors. |
| Importance | Coronary heart disease (CHD) is the most common cause of death in adults in the United States. Treatment to prevent CHD events by modifying risk factors is currently based on the Framingham risk model. If the classification of individuals at intermediate risk could be improved by using additional risk factors, treatment to prevent CHD might be targeted more effectively. Risk factors not currently part of the Framingham model (nontraditional risk factors) include high sensitivity C-reactive protein (hs-CRP), ankle-brachial index (ABI), leukocyte count, fasting blood glucose level, periodontal disease, carotid intima-media thickness, electron beam computed tomography, homocysteine level, and lipoprotein(a) level. |
| Balacne of Benefits and Harms | There is insufficient evidence to determine the percentage of intermediate-risk individuals who would be reclassified by screening with nontraditional risk factors, other than hs-CRP and ABI. For individuals reclassified as high-risk on the basis of hs-CRP or ABI scores, data are not available to determine whether they benefit from additional treatments. Little evidence is available to determine the harms of using nontraditional risk factors in screening. Potential harms include lifelong use of medications without proven benefit and psychological and other harms from being misclassified in a higher risk category. |
| Suggestions for practice | Clinicians should continue to use the Framingham model to assess CHD risk and guide risk-based preventive therapy. Adding nontraditional risk factors to CHD assessment would require additional patient and clinical staff time and effort. Routinely screening with nontraditional risk factors could result in lost opportunities to provide other important health services of proven benefit. |
| Other Relevant USPSTF Recommendations | USPSTF recommendations on risk assessment for CHD, the use of aspirin to prevent cardiovascular disease, and screening for high blood pressure can be accessed at http://www.uspreventiveservicestaskforce.org . |

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Coronary Heart Disease (Electrocardiography)

| Screening for Coronary Heart Disease with Electrocardiography | |
|--|---|
| Population | Asymptomatic adults at low risk for coronary heart disease (CHD) events |
| Recommendation | Asymptomatic adults at intermediate or high risk for CHD events No recommendation. Grade: I (Insufficient Evidence) |
| Risk Assessment | <p>Several factors are associated with a higher risk for CHD events, including older age, male sex, high blood pressure, smoking, abnormal lipid levels, diabetes, obesity, and sedentary lifestyle. Calculators are available to ascertain a person's 10-year risk for a CHD event.</p> <p>Persons with a 10-year risk >20% are considered to be high-risk, those with a 10-year risk <10% are considered to be low-risk, and those in the 10%–20% range are considered to be intermediate-risk.</p> |
| Screening Tests | <p>Several abnormalities on resting and exercise ECG are associated with an increased risk for a serious CHD event. However, the incremental information offered by screening asymptomatic adults at low risk for a CHD event with resting or exercise ECG (beyond that obtained with conventional CHD risk factors) is highly unlikely to result in changes in risk stratification that would prompt interventions and ultimately reduce CHD-related events.</p> |
| Balance of Benefits and Harms | <p>The potential harms of screening for CHD with exercise or resting ECG equal or exceed the potential benefits in this population.</p> <p>The USPSTF could not determine the balance between the benefits and harms of screening for CHD with resting or exercise ECG in this population.</p> |
| Other Relevant USPSTF Recommendations | <p>The USPSTF has made recommendations on screening for carotid artery stenosis, high blood pressure, lipid disorders, peripheral arterial disease, and obesity. These recommendations are available at http://www.uspreventiveservicestaskforce.org/.</p> |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Depression in Adults

| Screening for Depression in Adults | |
|--|---|
| Title | Screening for Depression in Adults |
| Population | Nonpregnant adults 18 years or older |
| Recommendation | <p style="text-align: center;">Screen when staff-assisted depression care supports¹ are in place to assure accurate diagnosis, effective treatment, and followup.</p> <p style="text-align: center;">Grade: B</p> <p style="text-align: center;">Do not automatically screen when staff-assisted depression care supports¹ are not in place.</p> <p style="text-align: center;">Grade: C</p> |
| Risk Assessment | Persons at increased risk for depression are considered at risk throughout their lifetime. Groups at increased risk include persons with other psychiatric disorders, including substance misuse; persons with a family history of depression; persons with chronic medical diseases; and persons who are unemployed or of lower socioeconomic status. Also, women are at increased risk compared with men. However, the presence of risk factors alone cannot distinguish depressed patients from nondepressed patients. |
| Screening Tests | Simple screening questions may perform as well as more complex instruments. Any positive screening test result should trigger a full diagnostic interview using standard diagnostic criteria. |
| Timing of Screening | The optimal interval for screening is unknown. In older adults, significant depressive symptoms are associated with common life events, including medical illness, cognitive decline, bereavement, and institutional placement in residential or inpatient settings. |
| Balance of Benefits and Harms | Limited evidence suggests that screening for depression in the absence of staff-assisted depression care does not improve depression outcomes. |
| Suggestions for Practice | "Staff-assisted depression care supports" refers to clinical staff that assists the primary care clinician by providing some direct depression care and/or coordination, case management, or mental health treatment. |
| Relevant USPSTF Recommendations | Related USPSTF recommendations on screening for suicidality and screening children and adolescents for depression are available at http://www.uspreventiveservicestaskforce.org . |

¹Go to the Suggestions for Practice section of this figure for further explanation.

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Diabetes Mellitus

| Screening for Type 2 Diabetes Mellitus in Adults | |
|--|--|
| Title | Screening for Type 2 Diabetes Mellitus in Adults |
| Population | Asymptomatic adults with sustained blood pressure greater than 135/80 mm Hg Asymptomatic adults with sustained blood pressure 135/80 mm Hg or lower |
| Recommendation | No recommendation. Grade: I (Insufficient Evidence) |
| Risk Assessment | These recommendations apply to adults with no symptoms of type 2 diabetes mellitus or evidence of possible complications of diabetes. Blood pressure measurement is an important predictor of cardiovascular complications in people with type 2 diabetes mellitus. The first step in applying this recommendation should be measurement of blood pressure (BP). Adults with treated or untreated BP >135/80 mm Hg should be screened for diabetes. |
| Screening Tests | Three tests have been used to screen for diabetes: <ul style="list-style-type: none"> ● Fasting plasma glucose (FPG). ● 2-hour postload plasma. ● Hemoglobin A1c. The American Diabetes Association (ADA) recommends screening with FPG, defines diabetes as FPG ≥ 126 mg/dL, and recommends confirmation with a repeated screening test on a separate day. |
| Screening Intervals | The optimal screening interval is not known. The ADA, on the basis of expert opinion, recommends an interval of every 3 years. |
| Suggestions for practice regarding insufficient evidence | When BP is ≤ 135/80 mm Hg, screening may be considered on an individual basis when knowledge of diabetes status would help inform decisions about coronary heart disease (CHD) preventive strategies, including consideration of lipid-lowering agents or aspirin. To determine whether screening would be helpful on an individual basis, information about 10-year CHD risk must be considered. For example, if CHD risk without diabetes was 17% and risk with diabetes was >20%, screening for diabetes would be helpful because diabetes status would determine lipid treatment. In contrast, if risk without diabetes was 10% and risk with diabetes was 15%, screening would not affect the decision to use lipid-lowering treatment. |
| Other relevant information from the USPSTF and the Community Preventive Services Task Force | Evidence and USPSTF recommendations regarding blood pressure, diet, physical activity, and obesity are available at http://www.uspreventiveservicestaskforce.org . The reviews and recommendations of the Community Preventive Services Task Force may be found at http://www.thecommunityguide.org . |

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Falls in Older Adults

| Title | Prevention of Falls in Community-Dwelling Older Adults | |
|--|--|---|
| Population | Community-dwelling adults aged 65 years and older who are at increased risk for falls | Community-dwelling adults aged 65 years and older |
| Recommendation | Provide intervention consisting of exercise or physical therapy and/or vitamin D supplementation to prevent falls. Grade: B | Do not automatically perform an in-depth multifactorial risk assessment with comprehensive management of identified risks to prevent falls. Grade: C |
| Risk Assessment | Primary care clinicians can consider the following factors to identify older adults at increased risk for falls: a history of falls, a history of mobility problems, and poor performance on the timed Get-Up-and-Go test. | |
| Interventions | <p>Effective exercise and physical therapy interventions include group classes and at-home physiotherapy strategies and range in intensity from very low (≤ 9 hours) to high (>75 hours).</p> <p>Benefit from vitamin D supplementation occurs by 12 months; the efficacy of treatment of shorter duration is unknown. The recommended daily allowance for vitamin D is 600 IU for adults aged 51 to 70 years and 800 IU for adults older than 70 years.</p> <p>Comprehensive multifactorial assessment and management interventions include assessment of multiple risk factors for falls and providing medical and social care to address factors identified during the assessment. In determining whether this service is appropriate in individual cases, patients and clinicians should consider the balance of benefits and harms on the basis of the circumstances of prior falls, medical comorbid conditions, and patient values.</p> | |
| Balance of Benefits and Harms | Exercise or physical therapy and vitamin D supplementation have a moderate benefit in preventing falls in older adults. | Multifactorial risk assessment with comprehensive management of identified risks has at least a small benefit in preventing falls in older adults. |
| Other Relevant USPSTF Recommendations | The USPSTF has made recommendations on screening for osteoporosis. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . | |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Folic Acid Supplementation

| | |
|-----------------------|--|
| Title | Folic Acid for the Prevention of Neural Tube Defects |
| Population | Women planning a pregnancy or capable of becoming pregnant |
| Recommendation | Take a daily vitamin supplement containing 0.4 to 0.8 mg (400 to 800 µg) of folic acid. Grade: A |

| | |
|----------------------------------|---|
| Risk Assessment | <p>Risk factors include:</p> <ul style="list-style-type: none"> ● A personal or family history of a pregnancy affected by a neural tube defect ● The use of certain antiseizure medications ● Mutations in folate-related enzymes ● Maternal diabetes ● Maternal obesity <p>Note: This recommendation does not apply to women who have had a previous pregnancy affected by neural tube defects or women taking certain antiseizure medicines. These women may be advised to take higher doses of folic acid.</p> |
| Timing of Medication | Start supplementation at least 1 month before conception. Continue through first 2 to 3 months of pregnancy. |
| Recommendations of Others | ACOG, AAFP, and most other organizations recommend 4 mg/d for women with a history of a pregnancy affected by a neural tube defect. |

Abbreviations: AAFP: American Academy of Family Physicians; ACOG: American College of Obstetricians and Gynecologists.

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Genital Herpes

| Screening for Genital Herpes | |
|--|--|
| Time | Asymptomatic adolescents and adults |
| Population | Asymptomatic pregnant women |
| Recommendation | Do not screen for herpes simplex virus. Grade: D |
| Screening Tests | Methods for detecting herpes simplex virus include viral culture, polymerase chain reaction, and antibody-based tests, such as the western blot assay and type-specific glycoprotein G serological tests. |
| Interventions | Antiviral therapy improves health outcomes in symptomatic persons (e.g., those with multiple recurrences); however, there is no evidence that the use of antiviral therapy improves health outcomes in those with asymptomatic infection. There are multiple efficacious regimens that may be used to prevent the recurrence of clinical genital herpes. |
| Balance of Benefits and Harms | <p>The potential harms of screening asymptomatic pregnant women include false-positive test results, labeling, and anxiety, as well as false-negative tests and false reassurance, although these potential harms are not well studied. The USPSTF determined that there are no benefits associated with screening, and therefore the potential harms outweigh the benefits.</p> <p>The potential harms of screening asymptomatic adolescents and adults include false-positive test results, labeling, and anxiety, although these potential harms are not well studied. The USPSTF determined the benefits of screening are minimal, at best, and the potential harms outweigh the potential benefits.</p> |
| Other Relevant USPSTF Recommendations | The USPSTF has made recommendations on screening for chlamydia, gonorrhea, HIV, and several other sexually transmitted infections. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Gestational Diabetes Mellitus

| Screening for Gestational Diabetes Mellitus | |
|--|--|
| Title | Screening for Gestational Diabetes Mellitus |
| Population | Asymptomatic pregnant women after 24 weeks of gestation |
| Recommendation | Asymptomatic pregnant women before 24 weeks of gestation No recommendation. Grade: I (Insufficient Evidence) |
| Risk Assessment | Risk factors that increase a woman's risk for developing GDM include obesity, increased maternal age, history of GDM, family history of diabetes, and belonging to an ethnic group with increased risk of developing type 2 diabetes mellitus (Hispanic, Native American, South or East Asian, African American, or Pacific Islands descent). There are 2 strategies used to screen for gestational diabetes in the United States. In the 2-step approach, the 50-g oral glucose challenge test is administered between 24 and 28 weeks of gestation in a nonfasting state. If the screening threshold is met or exceeded (7.22 mmol/L [130 mg/dL], 7.50 mmol/L [135 mg/dL], or 7.77 mmol/L [140 mg/dL]), patients receive the oral glucose tolerance test. A diagnosis of GDM is made when 2 or more glucose levels meet or exceed the specified glucose thresholds. In the 1-step approach, a 75-g glucose load is administered after fasting and plasma glucose levels are evaluated after 1 and 2 hours. GDM is diagnosed if 1 glucose value falls at or above the specified glucose threshold. Other methods of screening include fasting plasma glucose and screening based on risk factors. However, there is limited evidence on these alternative screening approaches. |
| Screening Tests | Initial treatment includes moderate physical activity, dietary changes, support from diabetes educators and nutritionists, and glucose monitoring. If the patient's glucose is not controlled after these initial interventions, she may be prescribed medication (either insulin or oral hypoglycemic agents), undergo increased surveillance in prenatal care, and have changes in delivery management. |
| Treatment | There is a moderate net benefit to screening for GDM after 24 weeks of gestation to reduce maternal and fetal complications. |
| Balance of Benefits and Harms | The USPSTF has made recommendations on screening for type 2 diabetes. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . |
| Other Relevant USPSTF Recommendations | |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Glaucoma

| | |
|--|---|
| Title | Screening for Glaucoma |
| Population | Adults without vision symptoms who are seen in primary care |
| Recommendation | No recommendation. Grade: I (Insufficient Evidence) |
| Risk Assessment | Important risk factors for open-angle glaucoma are increased intraocular pressure, older age, family history of glaucoma, and African American race. |
| Screening Tests | Diagnosis of glaucoma is usually made on the basis of several tests that, when combined, evaluate the biologic structure and function of the optic nerve and intraocular pressure. Most tests that are available in a primary care setting do not have acceptable accuracy to detect glaucoma. |
| Treatment | The immediate physiologic goal and measure of effect of primary treatment of glaucoma is reduction in intraocular pressure. Treatments that are effective in reducing intraocular pressure include medications, laser therapy, and surgery. However, these treatments have potential harms, and their effectiveness in reducing patient-perceived impairment in vision-related function is uncertain. |
| Balance of Benefits and Harms | Evidence on the accuracy of screening tests, especially in primary care settings, and the benefits of screening or treatment to delay or prevent visual impairment or improve quality of life is inadequate. Therefore, the overall certainty of the evidence is low, and the USPSTF is unable to determine the balance of benefits and harms of screening for glaucoma in asymptomatic adults. |
| Other Relevant USPSTF Recommendations | The USPSTF has made recommendations on screening for impaired visual acuity in older adults. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Gonorrhea

| Screening for Gonorrhea | | | |
|-------------------------|--|--|--|
| Title | Screening for Gonorrhea | | |
| Population | Sexually active women, including those who are pregnant, who are at increased risk for infection | Men who are at increased risk for infection | Men and women who are at low risk for infection |
| Recommendation | Screen for gonorrhea. Grade: B | No recommendation. Grade: I (Insufficient Evidence) | Do not screen for gonorrhea. Grade: D |
| | | No recommendation. Grade: I (Insufficient Evidence) | No recommendation. Grade: I (Insufficient Evidence) |

| | | | |
|--|--|---|--|
| Risk Assessment | Women and men younger than age 25 years—including sexually active adolescents—are at highest risk for gonorrhea infection. Risk factors for gonorrhea include a history of previous gonorrhea infection, other sexually transmitted infections, new or multiple sexual partners, inconsistent condom use, sex work, and drug use. Risk factors for pregnant women are the same as for non-pregnant women. | | |
| Screening Tests | Vaginal culture is an accurate screening test when transport conditions are suitable. Newer screening tests, including nucleic acid amplification and hybridization tests, have demonstrated improved sensitivity and comparable specificity when compared with cervical culture. Some newer tests can be used with urine and vaginal swabs, which enables screening when a pelvic examination is not performed. | | |
| Timing of Screening | Screening is recommended at the first prenatal visit for pregnant women who are in a high-risk group for gonorrhea infection. For pregnant women who are at continued risk, and for those who acquire a new risk factor, a second screening should be conducted during the third trimester. The optimal interval for screening in the non-pregnant population is not known. | | |
| Interventions | Genital gonorrhea infection in men and women, including pregnant women, may be treated with a third-generation cephalosporin. Because of increased prevalence of resistant organisms, fluoroquinolones should not be used to treat gonorrhea. Current guidelines for treating gonorrhea infection are available from the Centers for Disease Control and Prevention (http://www.cdc.gov/std/treatment). | | |
| Balance of Benefits and Harms | The USPSTF concluded that the benefits of screening women at increased risk for gonorrhea infection outweigh the potential harms. | The USPSTF could not determine the balance of benefits and harms of screening for gonorrhea in men at increased risk for infection. | Given the low prevalence of gonorrhea infection in the general population, the USPSTF concluded that the potential harms of screening in low-prevalence populations outweigh the benefits. |
| Other Relevant USPSTF Recommendations | The USPSTF has also made a recommendation on ocular prophylaxis in newborns for gonococcal ophthalmia neonatorum. This recommendation is available at http://www.uspreventiveservicestaskforce.org/ . | | |
| | The USPSTF could not determine the balance between the benefits and harms of screening for gonorrhea in pregnant women who are not at increased risk for infection. | | |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Healthful Diet and Physical Activity

| | |
|-----------------------|---|
| Title | Behavioral Counseling Interventions to Promote A Healthful Diet and Physical Activity for Cardiovascular Disease Prevention in Adults |
| Population | General adult population without a known diagnosis of hypertension, diabetes, hyperlipidemia, or cardiovascular disease |
| Recommendation | <p>Although the correlation among healthful diet, physical activity, and the incidence of cardiovascular disease is strong, existing evidence indicates that the health benefit of initiating behavioral counseling in the primary care setting to promote a healthful diet and physical activity is small. Clinicians may choose to selectively counsel patients rather than incorporate counseling into the care of all adults in the general population.</p> <p>Considerations: Issues to consider include other risk factors for cardiovascular disease, patient readiness for change, social support and community resources that support behavioral change, and other health care and preventive service priorities.</p> <p>Potential Harms: Harms may include the lost opportunity to provide other services with a greater health effect.</p> <p>Grade: C</p> |

| | |
|--|---|
| Risk Assessment | If an individual's risk for cardiovascular disease is uncertain, there are several calculators and models available to quantify a person's 10-year risk for cardiac events, such as the Framingham-based Adult Treatment Panel III calculator (available at http://hp2010.nhlbi.nih.gov/atpiii/calculator.asp). Generally, persons with a 10-year risk greater than 20% are considered to be high-risk, those with a 10-year risk less than 10% are considered to be low-risk, and those in the 10% to 20% range are considered to be intermediate-risk. |
| Interventions | Medium- or high-intensity behavioral interventions to promote a healthful diet and physical activity may be provided to individual patients in primary care settings or in other sectors of the health care system after referral from a primary care clinician. In addition, clinicians may offer healthful diet and physical activity interventions by referring the patient to community-based organizations. Strong linkages between the primary care setting and community-based resources may improve the delivery of these services. |
| Balance of Benefits and Harms | The USPSTF concludes with moderate certainty that medium- or high-intensity primary care behavioral counseling interventions to promote a healthful diet and physical activity have a small net benefit in adult patients without cardiovascular disease, hypertension, hyperlipidemia, or diabetes. |
| Other Relevant USPSTF Recommendations | The USPSTF has made recommendations on screening for carotid artery stenosis, coronary heart disease, high blood pressure, lipid disorders, peripheral arterial disease, and obesity. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Hearing Loss in Older Adults

| | |
|--------------------------------------|---|
| Title | Screening for Hearing Loss in Older Adults |
| Population | Asymptomatic adults aged 50 years or older |
| Recommendation | No recommendation. Grade: I (Insufficient Evidence) |
| Risk Assessment | Increasing age is the most important risk factor for hearing loss. Other risk factors include a history of exposure to loud noises or ototoxic agents, including occupational exposures, previous recurrent inner ear infections, genetic factors, and certain systemic diseases, such as diabetes. Various screening tests are used in primary care settings to detect hearing loss in adults, including: <ul style="list-style-type: none"> ● Whispered voice test ● Finger rub test ● Watch tick test ● Single-item screening (for example, asking “Do you have difficulty with your hearing?”) ● Multiple-item patient questionnaire (for example, Hearing Handicap Inventory for the Elderly—Screening Version) ● Handheld audiometer |
| Screening Tests | |
| Interventions | Hearing aids can improve self-reported hearing, communication, and social functioning for some adults with age-related hearing loss. |
| Balance of Benefits and Harms | There is inadequate evidence to determine the balance of benefits and harms of screening for hearing loss in adults aged 50 years or older. |

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Hemochromatosis

| | |
|--|---|
| Title | Screening for Hemochromatosis |
| Population | Asymptomatic general population |
| Recommendation | Do not screen for hereditary hemochromatosis. Grade: D |
| Risk Assessment | Clinically recognized hereditary hemochromatosis is primarily associated with mutations on the hemochromatosis (<i>HFE</i>) gene. Although this is a relatively common mutation in the U.S. population, only a small subset will develop symptoms of hemochromatosis. An even smaller proportion of these individuals will develop advanced stages of clinical disease. |
| Screening Tests | Genetic screening for <i>HFE</i> mutations can accurately identify individuals at risk for hereditary hemochromatosis. However, identifying an individual with the genotypic predisposition does not accurately predict the future risk for disease manifestation. |
| Interventions | Therapeutic phlebotomy is the main treatment for hereditary hemochromatosis. Phlebotomy is generally thought to have few side effects. |
| Balance of Benefits and Harms | <ul style="list-style-type: none"> • Screening could lead to identification of a large number of individuals who possess the high-risk genotype but may never manifest the clinical disease. This may result in unnecessary surveillance and diagnostic procedures, labeling, anxiety, and, potentially, unnecessary treatments. • There is poor evidence that early therapeutic phlebotomy improves morbidity and mortality in individuals with screening-detected versus clinically-detected hemochromatosis. • The USPSTF concluded that the potential harms of genetic screening for hereditary hemochromatosis outweigh the potential benefits. |
| Other Relevant USPSTF Recommendations | The USPSTF has also made recommendations on genetic testing for mutations in the breast cancer susceptibility gene to predict breast and ovarian cancer susceptibility. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Hepatitis B Virus Infection (Pregnant Women)

| | |
|--|--|
| Title | Screening for Hepatitis B Virus Infection in Pregnancy |
| Population | All pregnant women |
| Recommendation | Screen for hepatitis B virus (HBV) at the first prenatal visit. Grade: A |
| Screening Tests | Serologic identification of hepatitis B surface antigen (HBsAg). Reported sensitivity and specificity are greater than 98%. |
| Timing of Screening | Order HBsAg testing at the first prenatal visit. Re-screen women with unknown HBsAg status or new or continuing risk factors at admission to hospital, birth center, or other delivery setting. |
| Interventions | Administer hepatitis B vaccine and hepatitis B immune globulin to HBV-exposed infants within 12 hours of birth. Refer women who test positive for counseling and medical management. Counseling should include information about how to prevent transmission to sexual partners and household contacts. Reassure patients that breastfeeding is safe for infants who receive appropriate prophylaxis. |
| Implementation | Establish systems for timely transfer of maternal HBsAg test results to the labor and delivery and newborn medical records. |
| Other Relevant USPSTF Recommendations | USPSTF recommendations on the screening of pregnant women for other infections, including asymptomatic bacteriuria, bacterial vaginosis, chlamydia, HIV, and syphilis, can be found at http://www.uspreventiveservicestaskforce.org . |

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Hepatitis C Virus Infection

| | |
|--|---|
| Title | Screening for Hepatitis C Virus Infection in Adults |
| Population | Persons at high risk for infection and adults born between 1945 and 1965 |
| Recommendation | Screen for hepatitis C virus (HCV) infection. Grade: B |
| Risk Assessment | The most important risk factor for HCV infection is past or current injection drug use. Additional risk factors include receiving a blood transfusion before 1992, long-term hemodialysis, being born to an HCV-infected mother, incarceration, intranasal drug use, getting an unregulated tattoo, and other percutaneous exposures. Adults born between 1945 and 1965 are more likely to be diagnosed with HCV infection, either because they received a blood transfusion before the introduction of screening in 1992 or because they have a history of other risk factors for exposure decades earlier. |
| Screening Tests | Anti-HCV antibody testing followed by confirmatory polymerase chain reaction testing accurately identifies patients with chronic HCV infection. Various noninvasive tests with good diagnostic accuracy are possible alternatives to liver biopsy for diagnosing fibrosis or cirrhosis. |
| Screening Interval | Persons with continued risk for HCV infection (such as injection drug users) should be screened periodically. Evidence on how often screening should occur in these persons is lacking. Adults born between 1945 and 1965 and persons who are at risk because of potential exposure before universal blood screening need only be screened once. |
| Treatment | Antiviral treatment prevents long-term health complications of HCV infection (such as cirrhosis, liver failure, and hepatocellular carcinoma). The combination of pegylated interferon (α -2a or α -2b) and ribavirin is the standard treatment for HCV infection. In 2011, the U.S. Food and Drug Administration approved the protease inhibitors boceprevir and telaprevir for the treatment of HCV genotype 1 infection (the predominant genotype in the United States). |
| Balance of Benefits and Harms | On the basis of the accuracy of HCV antibody testing and the availability of effective interventions for persons with HCV infection, the USPSTF concludes that there is a moderate net benefit to screening in populations at high risk for infection. The USPSTF concludes that there is also a moderate net benefit to 1-time screening in all adults in the United States born between 1945 and 1965. |
| Other Relevant USPSTF Recommendations | The USPSTF has made recommendations on screening for hepatitis B virus infection in adolescents, adults, and pregnant women. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

High Blood Pressure in Adults

| Title | Screening for High Blood Pressure in adults |
|---------------------------------------|--|
| Population | Adult general population ¹ |
| Recommendation | Screen for high blood pressure. Grade: A |
| Screening Tests | High blood pressure (hypertension) is usually defined in adults as: systolic blood pressure (SBP) of 140 mm Hg or higher, or diastolic blood pressure (DBP) of 90 mm Hg or higher. Due to variability in individual blood pressure measurements, it is recommended that hypertension be diagnosed only after 2 or more elevated readings are obtained on at least 2 visits over a period of 1 to several weeks. |
| Screening Intervals | The optimal interval for screening adults for hypertension is not known. The Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC 7) recommends: <ul style="list-style-type: none"> ● Screening every 2 years with BP <120/80. ● Screening every year with SBP of 120-139 mmHg or DBP of 80-90 mmHg. |
| Treatment | A variety of pharmacological agents are available to treat hypertension. JNC 7 guidelines for treatment of hypertension can be accessed at http://www.nhlbi.nih.gov/guidelines/hypertension/jncintro.htm . The following non-pharmacological therapies are associated with reductions in blood pressure: <ul style="list-style-type: none"> ● Reduction of dietary sodium intake. ● Potassium supplementation. ● Increased physical activity, weight loss. ● Stress management. ● Reduction of alcohol intake. |
| Other Relevant USPSTF Recommendations | Adults with hypertension should be screened for diabetes. Adults should be screened for hyperlipidemia (depending on age, sex, risk factors) and smoking. Clinicians should discuss aspirin chemoprevention with patients at increased risk for cardiovascular disease. These recommendations and related evidence are available at http://www.uspreventiveservicestaskforce.org . |

¹This recommendation applies to adults without known hypertension.

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

HIV Infection

| | |
|--|---|
| Title | Screening for HIV |
| Population | Adolescents and adults aged 15 to 65 years, younger adolescents and older adults at increased risk for infection, and pregnant women |
| Recommendation | Screen for HIV infection. Grade: A |
| Risk Assessment | <p>Men who have sex with men and active injection drug users are at very high risk for new HIV infection. Other persons at high risk include those who have acquired or request testing for other sexually transmitted infections.</p> <p>Behavioral risk factors for HIV infection include:</p> <ul style="list-style-type: none"> ● Having unprotected vaginal or anal intercourse ● Having sexual partners who are HIV-infected, bisexual, or injection drug users ● Exchanging sex for drugs or money <p>The USPSTF recognizes that the above categories are not mutually exclusive, the degree of sexual risk is on a continuum, and individuals may not be aware of their sexual partners' risk factors for HIV infection.</p> |
| Screening Tests | <p>The conventional serum test for diagnosing HIV infection is repeatedly reactive immunoassay, followed by confirmatory Western blot or immunofluorescent assay. Conventional HIV test results are available within 1 to 2 days from most commercial laboratories.</p> <p>Rapid HIV testing may use either blood or oral fluid specimens and can provide results in 5 to 40 minutes; however, initial positive results require confirmation with conventional methods.</p> <p>Other U.S. Food and Drug Administration–approved tests for detection and confirmation of HIV infection include combination tests (for p24 antigen and HIV antibodies) and qualitative HIV-1 RNA.</p> |
| Interventions | <p>At present, there is no cure for chronic HIV infection. However, appropriately timed interventions in HIV-positive persons can reduce risks for clinical progression, complications or death from the disease, and disease transmission. Effective interventions include antiretroviral therapy (ART) (specifically, the use of combined ART), immunizations, and prophylaxis for opportunistic infections.</p> |
| Balance of Benefits and Harms | <p>The net benefit of screening for HIV infection in adolescents, adults, and pregnant women is substantial.</p> |
| Other Relevant USPSTF Recommendations | <p>The USPSTF has made recommendations on behavioral counseling to prevent sexually transmitted infections. This recommendation is available at http://www.uspreventiveservicestaskforce.org/.</p> |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Illicit Drug Use

| | |
|--|--|
| Title | Screening for Illicit Drug Use |
| Population | Adolescents ¹ , adults, and pregnant women not previously identified as users of illicit drugs |
| Recommendation | No recommendation. Grade I: (Insufficient Evidence) |
| Screening Tests | Toxicologic tests of blood or urine can provide objective evidence of drug use, but do not distinguish occasional users from impaired drug users. Valid and reliable standardized questionnaires are available to screen adolescents and adults for drug use or misuse. There is insufficient evidence to evaluate the clinical utility of these instruments when widely applied in primary care settings. |
| Balance of Benefits and Harms | The USPSTF concludes that for adolescents, adults, and pregnant women, the evidence is insufficient to determine the benefits and harms of screening for illicit drug use. |
| Suggestions for Practice | Clinicians should be alert to the signs and symptoms of illicit drug use in patients. |
| Treatment | More evidence is needed on the effectiveness of primary care office-based treatments for illicit drug use/dependence. |
| Other Relevant USPSTF Recommendations | The USPSTF recommendation for screening and counseling interventions to reduce alcohol misuse by adults and pregnant women can be found at http://www.uspreventiveservicestaskforce.org/uspstf/uspdrin.htm . |

¹For adolescents, see also Illicit and Prescription Drug Use in Children and Adolescents, Counseling

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Impaired Visual Acuity in Older Adults¹

| | |
|--|---|
| Title | Screening for Impaired Visual Acuity in Older Adults¹ |
| Population | Adults age 65 and older |
| Recommendation | No recommendation. Grade: I (Insufficient Evidence) |
| Risk Assessment | Older age is an important risk factor for most types of visual impairment. Additional risk factors include: <ul style="list-style-type: none"> • Smoking, alcohol use, exposure to ultraviolet light, diabetes, corticosteroids, and black race (for cataracts). • Smoking, family history, and white race (for age-related macular degeneration). |
| Screening Tests | Visual acuity testing (for example, the Snellen eye chart) is the usual method for screening for impairment of visual acuity in the primary care setting. Screening questions are not as accurate as a visual acuity test. |
| Balance of Benefits and Harms | There is no direct evidence that screening for vision impairment in older adults in primary care settings is associated with improved clinical outcomes. There is evidence that early treatment of refractive error, cataracts, and age-related macular degeneration may lead to harms that are small. The magnitude of net benefit for screening cannot be calculated because of a lack of evidence. |
| Other Relevant USPSTF Recommendations | Recommendations on screening for glaucoma and on screening for hearing loss in older adults can be accessed at http://www.uspreventiveservicestaskforce.org . |

¹This recommendation does not cover screening for glaucoma.

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Intimate Partner Violence and Elderly Abuse

| | | |
|-----------------------|--|--|
| Title | Screening for Intimate Partner Violence and Abuse of Elderly and Vulnerable Adults | |
| Population | Asymptomatic women of childbearing age | Elderly or vulnerable adults |
| Recommendation | Screen women for intimate partner violence (IPV), and provide or refer women who screen positive to intervention services. Grade: B | No recommendation. Grade: I (Insufficient Evidence) |

| | | |
|--|---|---|
| Risk Assessment | While all women are at potential risk for abuse, factors that elevate risk include young age, substance abuse, marital difficulties, and economic hardships. | |
| Interventions | Adequate evidence from randomized trials support a variety of interventions for women of childbearing age that can be delivered or referred by primary care, including counseling, home visits, information cards, referrals to community services, and mentoring support. Depending on the type of intervention, these services may be provided by clinicians, nurses, social workers, nonclinician mentors, or community workers. | |
| Balance of Benefits and Harms | Screening and interventions for IPV in women of childbearing age are associated with moderate health improvements through the reduction of exposure to abuse, physical and mental harms, and mortality. The associated harms are deemed no greater than small. Therefore, the overall net benefit is moderate. | The USPSTF was not able to estimate the magnitude of net benefit for screening all elderly or vulnerable adults (i.e., adults who are physically or mentally dysfunctional) for abuse and neglect because there were no studies on the accuracy, effectiveness, or harms of screening in this population. |
| Other Relevant USPSTF Recommendations | The USPSTF has made recommendations on screening for depression in adults and screening and counseling to reduce alcohol misuse in adults. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . | |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Lipid Disorders in Adults

| Title | Screening for Lipid Disorders in Adults | |
|-----------------------|---|--|
| Population | <ul style="list-style-type: none"> • Men age 35 years and older • Women age 45 years and older who are at increased risk for coronary heart disease (CHD) | <ul style="list-style-type: none"> • Men ages 20 to 35 years who are at increased risk for CHD • Women ages 20 to 45 years who are at increased risk for CHD |
| Recommendation | Screen for lipid disorders. Grade: A | Screen for lipid disorders. Grade: B |
| | | No recommendation for or against screening Grade: C |

| | | | |
|--|--|---|--|
| Risk Assessment | Consideration of lipid levels along with other risk factors allows for an accurate estimation of CHD risk. Risk factors for CHD include diabetes, history of previous CHD or atherosclerosis, family history of cardiovascular disease, tobacco use, hypertension, and obesity (body mass index ≥ 30 kg/m ²). | | |
| Screening Tests | The preferred screening tests for dyslipidemia are measuring serum lipid (total cholesterol, high-density and low-density lipoprotein cholesterol) levels in non-fasting or fasting samples. Abnormal screening results should be confirmed by a repeated sample on a separate occasion, and the average of both results should be used for risk assessment. | | |
| Timing of Screening | The optimal interval for screening is uncertain. Reasonable options include every 5 years, shorter intervals for people who have lipid levels close to those warranting therapy, and longer intervals for those not at increased risk who have had repeatedly normal lipid levels. An age at which to stop screening has not been established. Screening may be appropriate in older people who have never been screened; repeated screening is less important in older people because lipid levels are less likely to increase after age 65 years. | | |
| Interventions | Drug therapy is usually more effective than diet alone in improving lipid profiles, but choice of treatment should consider overall risk, costs of treatment, and patient preferences. Guidelines for treating lipid disorders are available from the National Cholesterol Education Program of the National Institutes of Health (http://www.nhlbi.nih.gov/about/ncep/). | | |
| Balance of Benefits and Harms | The benefits of screening for and treating lipid disorders in men age 35 and older and women age 45 and older at increased risk for CHD substantially outweigh the potential harms. | The benefits of screening for and treating lipid disorders in young adults at increased risk for CHD moderately outweigh the potential harms. | The net benefits of screening for lipid disorders in young adults not at increased risk for CHD are not sufficient to make a general recommendation. |
| Other Relevant USPSTF Recommendations | The USPSTF has made recommendations on screening for lipid disorders in children and screening for carotid artery stenosis, coronary heart disease, high blood pressure, and peripheral arterial disease. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . | | |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Lung Cancer

| | |
|--|---|
| Title | Screening for Lung Cancer |
| Population | Asymptomatic adults aged 55 to 80 years who have a 30 pack-year smoking history and currently smoke or have quit smoking within the past 15 years |
| Recommendation | Screen annually for lung cancer with low-dose computed tomography. Discontinue screening when the patient has not smoked for 15 years. Grade: B |
| Risk Assessment | Age, total cumulative exposure to tobacco smoke, and years since quitting smoking are the most important risk factors for lung cancer. Other risk factors include specific occupational exposures, radon exposure, family history, and history of pulmonary fibrosis or chronic obstructive lung disease. |
| Screening Tests | Low-dose computed tomography has high sensitivity and acceptable specificity for detecting lung cancer in high-risk persons and is the only currently recommended screening test for lung cancer. |
| Treatment | Non-small cell lung cancer is treated with surgical resection when possible and also with radiation and chemotherapy. |
| Balance of Benefits and Harms | Annual screening for lung cancer with low-dose computed tomography is of moderate net benefit in asymptomatic persons who are at high risk for lung cancer based on age, total cumulative exposure to tobacco smoke, and years since quitting smoking. |
| Other Relevant USPSTF Recommendations | The USPSTF has made recommendations on counseling and interventions to prevent tobacco use and tobacco-caused disease. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Menopausal Hormone Therapy

| Menopausal Hormone Therapy for the Primary Prevention of Chronic Conditions | |
|---|---|
| Title | Menopausal hormone therapy for the primary prevention of chronic conditions |
| Population | Postmenopausal women who have had a hysterectomy |
| Recommendation | <p>Do not prescribe combined estrogen and progestin for the prevention of chronic conditions.</p> <p>Grade: D</p> |

| | |
|---|---|
| <p>Risk Assessment</p> | <p>This recommendation applies to the average-risk population. Risk factors for a specific chronic disease or individual characteristics that affect the likelihood of a specific therapy-associated adverse event may cause a woman's net balance of benefits and harms to differ from that of the average population.</p> |
| <p>Preventive Medications</p> | <p>Although combined estrogen and progestin therapy (specifically, oral conjugated equine estrogen, 0.625 mg/d, plus medroxyprogesterone acetate, 2.5 mg/d) decreases the risk for fractures in postmenopausal women, there is an accompanying increased risk for serious adverse events, such as stroke, invasive breast cancer, dementia, gallbladder disease, deep venous thrombosis, and pulmonary embolism.</p> <p>Estrogen therapy (specifically, oral conjugated equine estrogen, 0.625 mg/d) decreases the risk for fractures and has a small effect on the risk for invasive breast cancer, but it is also associated with important harms, such as an increased likelihood of stroke, deep venous thrombosis, and gallbladder disease.</p> <p>Neither combined estrogen and progestin therapy nor estrogen alone reduces the risk for coronary heart disease in postmenopausal women.</p> |
| <p>Balance of Benefits and Harms</p> | <p>The chronic disease prevention benefits of combined estrogen and progestin do not outweigh the harms in most postmenopausal women.</p> <p>The chronic disease prevention benefits of estrogen are unlikely to outweigh the harms in most postmenopausal women who have had a hysterectomy.</p> |
| <p>Other Relevant USPSTF Recommendations</p> | <p>The USPSTF has made recommendations on screening for osteoporosis and the use of preventive medications for breast cancer, as well as other relevant interventions for the primary or secondary prevention of chronic diseases in women, such as medications for cardiovascular disease and screening for coronary heart disease, high blood pressure, lipid disorders, colorectal cancer, breast cancer, and dementia. These recommendations are available at http://www.uspreventiveservicestaskforce.org/.</p> |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Motor Vehicle Occupant Restraints

| | |
|---|--|
| Title | Primary Care Counseling for Proper Use of Motor Vehicle Occupant Restraints |
| Population | General primary care population |
| Recommendation | No recommendation. Grade: I (Insufficient Evidence) |
| Interventions | There is good evidence that community and public health interventions, including legislation, law enforcement campaigns, car seat distribution campaigns, media campaigns, and other community-based interventions, are effective in improving the proper use of car seats, booster seats, and seat belts. |
| Suggestions for Practice | Current evidence is insufficient to assess the incremental benefit of counseling in primary care settings, beyond increases related to other interventions, in improving rates of proper use of motor vehicle occupant restraints. Linkages between primary care and community interventions are critical for improving proper car seat, booster seat, and seat belt use. |
| Relevant Recommendations from the Guide to Community Preventive Services | The Community Preventive Services Task Force has reviewed evidence of the effectiveness of selected population-based interventions to reduce motor vehicle occupant injuries, focusing on three strategic areas: <ul style="list-style-type: none"> • Increasing the proper use of child safety seats. • Increasing the use of safety belts. • Reducing alcohol-impaired driving. Multiple interventions in these areas have been recommended. Recommendations can be accessed at http://www.thecommunityguide.org/mvoi/ |

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Obesity in Adults

| | |
|-----------------------|---|
| Title | Screening for and Management Of Obesity In Adults |
| Population | Adults aged 18 years or older |
| Recommendation | Screen for obesity. Patients with a body mass index (BMI) of 30 kg/m² or higher should be offered or referred to intensive, multicomponent behavioral interventions. Grade: B |

| | |
|--|---|
| Screening Tests | Body mass index is calculated from the measured weight and height of an individual. Recent evidence suggests that waist circumference may be an acceptable alternative to BMI measurement in some patient subpopulations. |
| Timing of Screening | No evidence was found about appropriate intervals for screening. |
| Interventions | Intensive, multicomponent behavioral interventions for obese adults include the following components: <ul style="list-style-type: none"> ● Behavioral management activities, such as setting weight-loss goals ● Improving diet or nutrition and increasing physical activity ● Addressing barriers to change ● Self-monitoring ● Strategizing how to maintain lifestyle changes |
| Balance of Benefits and Harms | Adequate evidence indicates that intensive, multicomponent behavioral interventions for obese adults can lead to weight loss, as well as improved glucose tolerance and other physiologic risk factors for cardiovascular disease. Inadequate evidence was found about the effectiveness of these interventions on long-term health outcomes (for example, mortality, cardiovascular disease, and hospitalizations). Adequate evidence indicates that the harms of screening and behavioral interventions for obesity are small. Possible harms of behavioral weight-loss interventions include decreased bone mineral density and increased fracture risk, serious injuries resulting from increased physical activity, and increased risk for eating disorders. |
| Other relevant USPSTF recommendations | Recommendations on screening for obesity in children and adolescents can be found at http://www.uspreventiveservicestaskforce.org/ . |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Oral Cancer

| | |
|--|--|
| Title | Screening for Oral Cancer |
| Population | Asymptomatic adults aged 18 years or older |
| Recommendation | No recommendation. Grade: I (Insufficient Evidence) |
| Risk Assessment | The primary risk factors for oral cancer are tobacco and alcohol use. Additional risk factors include male sex, older age, use of betel quid, ultraviolet light exposure, infection with <i>Candida</i> or bacterial flora, and a compromised immune system. Recently, sexually transmitted oral human papillomavirus infection has been recognized as an increasing risk factor for oropharyngeal cancer, another subset of head and neck cancer. |
| Screening Tests | The primary screening test for oral cancer is a systematic clinical examination, including inspection and palpation of the oral cavity. |
| Treatment | Suspected oral cancer or its precursors detected on examination require confirmation by tissue biopsy. Treatment for screen-detected oral cancer includes surgery, radiotherapy, and chemotherapy. |
| Balance of Benefits and Harms | The USPSTF found inadequate evidence on the diagnostic accuracy, benefits, and harms of screening for oral cancer. Therefore, the USPSTF cannot determine the balance of benefits and harms of screening for oral cancer in asymptomatic adults. |
| Other Relevant USPSTF Recommendations | The USPSTF has made recommendations on counseling to prevent tobacco use and screening for and counseling to reduce alcohol misuse. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Osteoporosis

| Screening for Osteoporosis | | |
|----------------------------|--|---|
| Title | Screening for Osteoporosis | |
| Population | Women age ≥65 years without previous known fractures or secondary causes of osteoporosis | Women age <65 years whose 10-year fracture risk is equal to or greater than that of a 65-year-old white woman without additional risk factors |
| Recommendation | Screen for osteoporosis. Grade: B | No recommendation. Grade: I (Insufficient Evidence) |

| | |
|---|---|
| Risk Assessment | As many as 1 in 2 postmenopausal women and 1 in 5 older men are at risk for an osteoporosis-related fracture. Osteoporosis is common in all racial groups but is most common in white persons. Rates of osteoporosis increase with age. Elderly people are particularly susceptible to fractures. According to the FRAX fracture risk assessment tool, available at http://www.shef.ac.uk/FRAX/ , the 10-year fracture risk in a 65-year-old white woman without additional risk factors is 9.3%. |
| Screening Tests | Current diagnostic and treatment criteria rely on dual-energy x-ray absorptiometry of the hip and lumbar spine. |
| Timing of Screening | Evidence is lacking about optimal intervals for repeated screening. |
| Intervention | In addition to adequate calcium and vitamin D intake and weight-bearing exercise, multiple U.S. Food and Drug Administration–approved therapies reduce fracture risk in women with low bone mineral density and no previous fractures, including bisphosphonates, parathyroid hormone, raloxifene, and estrogen. The choice of treatment should take into account the patient’s clinical situation and the tradeoff between benefits and harms. Clinicians should provide education about how to minimize drug side effects. |
| Suggestions for Practice Regarding the I Statement for Men | <p>Clinicians should consider:</p> <ul style="list-style-type: none"> ● potential preventable burden: increasing because of the aging of the U.S. population ● potential harms: likely to be small, mostly opportunity costs ● current practice: routine screening of men not widespread ● costs: additional scanners required to screen sizeable populations <p>Men most likely to benefit from screening have a 10-year risk for osteoporotic fracture equal to or greater than that of a 65-year-old white woman without risk factors. However, current evidence is insufficient to assess the balance of benefits and harms of screening for osteoporosis in men.</p> |

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Ovarian Cancer

| | |
|--|--|
| Title | Screening for Ovarian Cancer |
| Population | Asymptomatic women without known genetic mutations that increase risk for ovarian cancer |
| Recommendation | Do not screen for ovarian cancer. Grade: D |
| Risk Assessment | <p>Women with <i>BRCA1</i> and <i>BRCA2</i> genetic mutations, the Lynch syndrome (hereditary nonpolyposis colon cancer), or a family history of ovarian cancer are at increased risk for ovarian cancer.</p> <p>Women with an increased-risk family history should be considered for genetic counseling to further evaluate their potential risks. "Increased-risk family history" generally means having 2 or more first- or second-degree relatives with a history of ovarian cancer or a combination of breast and ovarian cancer; for women of Ashkenazi Jewish descent, it means having a first-degree relative (or 2 second-degree relatives on the same side of the family) with breast or ovarian cancer.</p> |
| Screening Tests | Transvaginal ultrasonography and serum cancer antigen (CA)-125 testing are the most commonly suggested screening modalities. |
| Treatments | Treatment of ovarian carcinoma includes surgical treatment (debulking) and intraperitoneal or systemic chemotherapy. |
| Balance of Benefits and Harms | Annual screening with transvaginal ultrasonography and serum CA-125 testing in women does not decrease ovarian cancer mortality. Screening for ovarian cancer can lead to important harms, including major surgical interventions in women who do not have cancer. Therefore, the harms of screening for ovarian cancer outweigh the benefits. |
| Other Relevant USPSTF Recommendations | The USPSTF has made a recommendation on genetic risk assessment and BRCA mutation testing for breast and ovarian cancer susceptibility. This recommendation is available at http://www.uspreventiveservicestaskforce.org/ . |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Peripheral Artery Disease

| | |
|--|--|
| Title | Screening for Peripheral Artery Disease and Cardiovascular Disease Risk Assessment with the Ankle Brachial Index in Adults |
| Population | Asymptomatic adults without a known diagnosis of peripheral artery disease (PAD), cardiovascular disease, severe chronic kidney disease, or diabetes |
| Recommendation | No recommendation. Grade: I (Insufficient Evidence) |
| Risk Assessment | Important risk factors for PAD include older age, diabetes, smoking, hypertension, high cholesterol level, obesity, and physical inactivity. Peripheral artery disease is more common in men than women and occurs at an earlier age in men. |
| Screening Tests | Resting ankle–brachial index (ABI) is the most commonly used test in screening for and detection of PAD in clinical settings. It is calculated as the systolic blood pressure obtained at the ankle divided by the systolic blood pressure obtained at the brachial artery while the patient is lying down. Physical examination has low sensitivity for detecting mild PAD in asymptomatic persons. |
| Balance of Benefits and Harms | Evidence on screening for PAD with the ABI in asymptomatic adults with no known diagnosis of cardiovascular disease or diabetes is insufficient; therefore, the balance of benefits and harms cannot be determined. |
| Other Relevant USPSTF Recommendations | The USPSTF has made recommendations on using nontraditional risk factors, including the ABI, in screening for coronary heart disease. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Prostate Cancer

| | |
|--|--|
| Title | Screening for Prostate Cancer |
| Population | Adult males |
| Recommendation | Do not use prostate-specific antigen (PSA)-based screening for prostate cancer. Grade: D |
| Screening Tests | <p>Contemporary recommendations for prostate cancer screening all incorporate the measurement of serum PSA levels; other methods of detection, such as digital rectal examination or ultrasonography, may be included.</p> <p>There is convincing evidence that PSA-based screening programs result in the detection of many cases of asymptomatic prostate cancer, and that a substantial percentage of men who have asymptomatic cancer detected by PSA screening have a tumor that either will not progress or will progress so slowly that it would have remained asymptomatic for the man's lifetime (i.e., PSA-based screening results in considerable overdiagnosis).</p> |
| Interventions | <p>Management strategies for localized prostate cancer include watchful waiting, active surveillance, surgery, and radiation therapy.</p> <p>There is no consensus regarding optimal treatment.</p> |
| Balance of Benefits and Harms | <p>The reduction in prostate cancer mortality 10 to 14 years after PSA-based screening is, at most, very small, even for men in the optimal age range of 55 to 69 years.</p> <p>The harms of screening include pain, fever, bleeding, infection, and transient urinary difficulties associated with prostate biopsy, psychological harm of false-positive test results, and overdiagnosis.</p> <p>Harms of treatment include erectile dysfunction, urinary incontinence, bowel dysfunction, and a small risk for premature death. Because of the current inability to reliably distinguish tumors that will remain indolent from those destined to be lethal, many men are being subjected to the harms of treatment for prostate cancer that will never become symptomatic.</p> <p>The benefits of PSA-based screening for prostate cancer do not outweigh the harms.</p> |
| Relevant USPSTF Recommendations | <p>Recommendations on screening for other types of cancer can be found at http://www.uspreventiveservicestaskforce.org/.</p> |

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Sexually Transmitted Infections

| Behavioral Counseling to Prevent Sexually Transmitted Infections | | | |
|--|--|--|---|
| Title | | | |
| Population | All sexually active adolescents | Adults at increased risk for STIs | Non-sexually-active adolescents and adults not at increased risk for STIs |
| Recommendation | Offer high-intensity counseling. Grade: B | Offer high-intensity counseling. Grade: B | No recommendation. Grade: I (Insufficient Evidence) |

| | |
|--|--|
| Risk Assessment | <p>All sexually active adolescents are at increased risk for STIs and should be offered counseling. Adults should be considered at increased risk and offered counseling if they have:</p> <ul style="list-style-type: none"> • Current STIs or have had an STI within the past year. • Multiple sexual partners. <p>In communities or populations with high rates of STIs, all sexually active patients in non-monogamous relationships may be considered at increased risk.</p> |
| Interventions | <p>Characteristics of successful high-intensity counseling interventions:</p> <ul style="list-style-type: none"> • Multiple sessions of counseling. • Frequently delivered in group settings. |
| Suggestions for Practice | <p>High-intensity counseling may be delivered in primary care settings, or in other sectors of the health system and community settings after referral.</p> <p>Delivery of this service may be greatly improved by strong linkages between the primary care setting and community.</p> <p>Evidence is limited regarding counseling for adolescents who are not sexually active. Intensive counseling for all adolescents in order to reach those who are at risk but have not been appropriately identified is not supported by current evidence. Evidence is lacking regarding the effectiveness of counseling for adults not at increased risk for STIs.</p> |
| Other Relevant USPSTF Recommendations | <p>USPSTF recommendations on screening for chlamydial infection, gonorrhea, genital herpes, hepatitis B, hepatitis C, HIV, and syphilis can be found at http://www.uspreventiveservicestaskforce.org.</p> |

Abbreviation: STI Sexually Transmitted Infection

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Skin Cancer (Counseling)

| Title | Behavioral Counseling to Prevent Skin Cancer | |
|--|--|---|
| Population | Children, adolescents, and young adults aged 10 to 24 years with fair skin | Adults older than age 24 years |
| Recommendation | <p style="text-align: center;">Provide counseling about minimizing exposure to ultraviolet radiation to reduce risk for skin cancer.</p> <p style="text-align: center;">Grade: B</p> | |
| Risk Assessment | <p>Individuals with a fair skin type are at greatly increased risk for skin cancer. Fair skin type can be defined by eye and hair color; freckling; and historical factors, such as usual reaction to sun exposure (always or usually burning or infrequently tanning).</p> | |
| Behavioral Counseling | <p>Effective counseling interventions were generally of low intensity and almost entirely accomplished within the primary care visit.</p> <p>Successful counseling interventions used cancer prevention or appearance-focused messages (such as stressing the aging effect of ultraviolet radiation on the skin) to reach specific audiences.</p> | |
| Interventions | <p>Behavior change interventions are aimed at reducing ultraviolet radiation exposure. Sun-protective behaviors include the use of a broad-spectrum sunscreen with a sun protection factor ≥ 15, wearing hats or other shade-protective clothing, avoiding the outdoors during midday hours (10 a.m. to 3 p.m.), and avoiding the use of indoor tanning.</p> | |
| Balance of Benefits and Harms | <p>For children, adolescents, and young adults aged 10 to 24 years with fair skin, primary care counseling interventions can increase the use of sun-protective behaviors by a moderate amount, with no appreciable harms.</p> | <p>For adults older than 24 years, there is inadequate evidence to determine the effect of counseling on the use of sun-protective behaviors.</p> |
| Other Relevant USPSTF Recommendations | <p>The USPSTF has made recommendations on screening for skin cancer. These recommendations are available at http://www.uspreventiveservicestaskforce.org/.</p> | |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Skin Cancer (Screening)

| Title | Screening for Skin Cancer |
|---|--|
| Population | Adult general population ¹ |
| Recommendation | No recommendation. Grade: I (Insufficient Evidence) |
| Risk Assessment | <p>Skin cancer risks: family history of skin cancer, considerable history of sun exposure and sunburn.</p> <p>Groups at increased risk for melanoma:</p> <ul style="list-style-type: none"> ● Fair-skinned men and women over the age of 65 years. ● Patients with atypical moles. ● Patients with more than 50 moles. |
| Screening Tests | There is insufficient evidence to assess the balance of benefits and harms of whole body skin examination by a clinician or patient skin self-examination for the early detection of skin cancer. |
| Screening Intervals | Not applicable. |
| Suggestions for Practice | Clinicians should remain alert for skin lesions with malignant features that are noted while performing physical examinations for other purposes. Features associated with increased risk for malignancy include: asymmetry, border irregularity, color variability, diameter >6mm ("A," "B," "C," "D"), or rapidly changing lesions. Suspicious lesions should be biopsied. |
| Other Relevant Recommendations from the USPSTF and the Community Preventive Services Task Force | <p>The USPSTF has reviewed the evidence for counseling to prevent skin cancer. The recommendation statement and supporting documents can be accessed at http://www.uspreventiveservicestaskforce.org.</p> <p>The Community Preventive Services Task Force has reviewed the evidence on public health interventions to reduce skin cancer. The recommendations can be accessed at http://www.thecommunityguide.org.</p> |

¹The USPSTF does not examine outcomes related to surveillance of patients with familial syndromes, such as familial atypical mole and melanoma (FAM-M) syndrome.

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Suicide Risk

| | |
|-----------------------|--|
| Title | Screening for Suicide Risk |
| Population | General population |
| Recommendation | No recommendation. Grade: I (Insufficient Evidence) |

| | |
|--|---|
| Risk Assessment | <p>The strongest risk factors for attempted suicide include mood disorders or other mental disorders, comorbid substance abuse disorders, history of deliberate self-harm, and a history of suicide attempts. Deliberate self-harm refers to intentionally initiated acts of self-harm with a nonfatal outcome (including self-poisoning and self-injury). Suicide risk is assessed along a continuum ranging from suicidal ideation alone (relatively less severe) to suicidal ideation with a plan (more severe). Suicidal ideation with a specific plan of action is associated with a significant risk for attempted suicide.</p> |
| Screening Tests | <p>There is limited evidence on the accuracy of screening tools to identify suicide risk in the primary care setting, including tools to identify those at high risk. The characteristics of the most commonly used screening instruments (Scale for Suicide Ideation, Scale for Suicide Ideation–Worst, and the Suicidal Ideation Questionnaire) have not been validated to assess suicide risk in primary care settings.</p> |
| Interventions | <p>There is insufficient evidence to determine if treatment of persons at high risk for suicide reduces suicide attempts or mortality.</p> |
| Balance of Benefits and Harms | <p>There is no evidence that screening for suicide risk reduces suicide attempts or mortality. There is insufficient evidence to determine if treatment of persons at high risk reduces suicide attempts or mortality. There are no studies that directly address the harms of screening and treatment for suicide risk.</p> <p>As a result, the USPSTF could not determine the balance of benefits and harms of screening for suicide risk in the primary care setting.</p> |
| Other Relevant USPSTF Recommendations | <p>The USPSTF has also made recommendations on screening for alcohol misuse, depression, and illicit drug use. These recommendations are available at http://www.uspreventiveservicestaskforce.org/.</p> |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Syphilis (Pregnant Women)

| | |
|--|---|
| Title | Screening for Syphilis Infection in Pregnancy |
| Population | All pregnant women |
| Recommendation | Screen for syphilis infection. Grade: A |
| Screening Tests | <p>Nontreponemal tests commonly used for initial screening include:</p> <ul style="list-style-type: none"> • Venereal Disease Research Laboratory (VDRL) • Rapid Plasma Reagin (RPR) <p>Confirmatory tests include:</p> <ul style="list-style-type: none"> • Fluorescent treponemal antibody absorbed (FTA-ABS) • <i>Treponema pallidum</i> particle agglutination (TPPA) |
| Timing of Screening | Test all pregnant women at the first prenatal visit. |
| Other Clinical Considerations | <p>Most organizations recommend testing high-risk women again during the third trimester and at delivery. Groups at increased risk include:</p> <ul style="list-style-type: none"> • Uninsured women • Women living in poverty • Sex workers • Illicit drug users • Those diagnosed with other sexually transmitted infections (STIs) • Other women living in communities with high syphilis morbidity <p>Prevalence is higher in southern U.S. and in metropolitan areas and in Hispanic and African American populations.</p> |
| Interventions | <p>The Centers for Disease Control and Prevention (CDC) recommends treatment with parenteral benzathine penicillin G. Women with penicillin allergies should be desensitized and treated with penicillin.</p> <p>Consult the CDC for the most up-to-date recommendations: http://www.cdc.gov/std/treatment/.</p> |
| Other Relevant USPSTF Recommendations | Recommendations on screening for other STIs, and on counseling for STIs, can be found at http://www.uspreventiveservicestaskforce.org . |

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Testicular Cancer

| | |
|--|--|
| Title | Screening for Testicular Cancer |
| Population | Adolescent and adult males |
| Recommendation | Do not screen. Grade: D |
| Screening Tests | There is inadequate evidence that screening asymptomatic patients by means of self-examination or clinician examination has greater yield or accuracy for detecting testicular cancer at more curable stages. |
| Interventions | Management of testicular cancer consists of orchiectomy and may include other surgery, radiation therapy, or chemotherapy, depending on stage and tumor type. Regardless of disease stage, over 90% of all newly diagnosed cases of testicular cancer will be cured. |
| Balance of Benefits and Harms | Screening by self-examination or clinician examination is unlikely to offer meaningful health benefits, given the very low incidence and high cure rate of even advanced testicular cancer. Potential harms include false-positive results, anxiety, and harms from diagnostic tests or procedures. |
| Other Relevant USPSTF Recommendations | Recommendations on screening for other types of cancer can be found at http://www.uspreventiveservicestaskforce.org . |

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Tobacco Use in Adults

| | | |
|-----------------------|--|--|
| Title | Counseling and Interventions to Prevent Tobacco Use and Tobacco-Caused Disease in Adults and Pregnant Women | |
| Population | Adults age ≥ 18 years | Pregnant women of any age |
| Recommendation | Ask about tobacco use. Provide tobacco cessation interventions to those who use tobacco products. | Ask about tobacco use. Provide augmented pregnancy-tailored counseling for women who smoke. |
| | Grade: A | Grade: A |

| | | |
|--|---|---|
| Counseling | <p>The “5-A” framework provides a useful counseling strategy:</p> <ol style="list-style-type: none"> 1. Ask about tobacco use. 2. Advise to quit through clear personalized messages. 3. Assess willingness to quit. 4. Assist to quit. 5. Arrange follow-up and support. <p>Intensity of counseling matters: brief one-time counseling works; however, longer sessions or multiple sessions are more effective.</p> <p>Telephone counseling “quit lines” also improve cessation rates.</p> | |
| Pharmacotherapy | <p>Combination therapy with counseling and medications is more effective than either component alone. FDA-approved pharmacotherapy includes nicotine replacement therapy, sustained-release bupropion, and varenicline.</p> | <p>The USPSTF found inadequate evidence to evaluate the safety or efficacy of pharmacotherapy during pregnancy.</p> |
| Implementation | <p>Successful implementation strategies for primary care practice include:</p> <ul style="list-style-type: none"> ● Instituting a tobacco user identification system. ● Promoting clinician intervention through education, resources, and feedback. ● Dedicating staff to provide treatment, and assessing the delivery of treatment in staff performance evaluations. | |
| Other Relevant USPSTF Recommendations | <p>Recommendations on other behavioral counseling topics are available at http://www.uspreventiveservicestaskforce.org.</p> | |

Abbreviations: FDA U.S. Food and Drug Administration; USPSTF U.S. Preventive Services Task Force

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Vitamin D and Calcium Supplementation to Prevent Fractures

| Vitamin D and Calcium Supplementation to Prevent Fractures in Adults | | | |
|--|--|---|---|
| Title | Vitamin D and Calcium Supplementation to Prevent Fractures in Adults | | |
| Population | Men or premenopausal women | Community-dwelling postmenopausal women at doses of >400 IU of vitamin D ₃ and >1,000 mg of calcium | Community-dwelling postmenopausal women at doses of ≤400 IU of vitamin D ₃ and ≤1,000 mg of calcium |
| Recommendation | No recommendation. Grade: I (Insufficient Evidence) | No recommendation. Grade: I (Insufficient Evidence) | Do not supplement. Grade: D recommendation |
| Preventive Medications | <p>Appropriate intake of vitamin D and calcium are essential to overall health. However, there is inadequate evidence to determine the effect of combined vitamin D and calcium supplementation on the incidence of fractures in men or premenopausal women.</p> <p>There is adequate evidence that daily supplementation with 400 IU of vitamin D₃ and 1,000 mg of calcium has no effect on the incidence of fractures in postmenopausal women.</p> <p>There is inadequate evidence regarding the effect of higher doses of combined vitamin D and calcium supplementation on fracture incidence in community-dwelling postmenopausal women.</p> | | |
| Balance of Benefits and Harms | Evidence is lacking regarding the benefit of daily vitamin D and calcium supplementation for the primary prevention of fractures, and the balance of benefits and harms cannot be determined. | Evidence is lacking regarding the benefit of daily supplementation with >400 IU of vitamin D ₃ and >1,000 mg of calcium for the primary prevention of fractures in postmenopausal women, and the balance of benefits and harms cannot be determined. | Daily supplementation with ≤400 IU of vitamin D ₃ and ≤1,000 mg of calcium has no net benefit for the primary prevention of fractures. |
| Other Relevant USPSTF Recommendations | The USPSTF has made recommendations on screening for osteoporosis and vitamin D supplementation to prevent falls in community-dwelling older adults. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . | | |


For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Vitamin Supplementation to Prevent Cardiovascular Disease and Cancer

| | | | |
|-----------------------|---|---|---|
| Title | Vitamin, Mineral, and Multivitamin Supplements for the Primary Prevention of Cardiovascular Disease and Cancer | | |
| Population | Healthy adults without special nutritional needs. This recommendation does not apply to children, women who are pregnant or may become pregnant, or persons who are chronically ill or hospitalized or have a known nutritional deficiency. | | |
| Recommendation | Multivitamins: No recommendation. Grade: I statement | Single- or paired-nutrient supplements: No recommendation. Grade: I statement | β-carotene or vitamin E: Do not recommend. Grade: D |

| | | | |
|--|---|--|---|
| Preventive Medications | Evidence on supplementation with multivitamins to reduce the risk for cardiovascular disease or cancer is inadequate, as is the evidence on supplementation with individual vitamins, minerals, or functional pairs. Supplementation with β-carotene or vitamin E does not reduce the risk for cardiovascular disease or cancer. | | |
| Balance of Benefits and Harms | The evidence is insufficient to determine the balance of benefits and harms of supplementation with multivitamins for the prevention of cardiovascular disease or cancer. | The evidence is insufficient to determine the balance of benefits and harms of supplementation with single or paired nutrients for the prevention of cardiovascular disease or cancer. | There is no net benefit of supplementation with vitamin E or β-carotene for the prevention of cardiovascular disease or cancer. |
| Other Relevant USPSTF Recommendations | The USPSTF has made several recommendations on the prevention of cardiovascular disease and cancer, including recommendations for smoking cessation, screening for lipid disorders, hypertension, diabetes, and cancer; obesity screening and counseling; and aspirin use. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . | | |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.



Clinical Summaries of Recommendations for Children and Adolescents

All clinical summaries in this Guide are abridged recommendations. To see the full recommendation statements and recommendations published after March 2014, go to www.USPreventiveServicesTaskForce.org.

Blood Lead Levels in Children and Pregnant Women

| Screening for Elevated Blood Lead Levels in Children and Pregnant Women | | | | |
|---|--|--|---|--|
| Population | Asymptomatic children ages 1 to 5 years who are at average risk Asymptomatic children ages 1 to 5 years who are at increased risk | | | |
| Recommendation | <p>No recommendation. Grade: I (Insufficient Evidence)</p> <p>Do not screen for elevated blood lead levels. Grade: D</p> <p>Do not screen for elevated blood lead levels. Grade: D</p> <p>Do not screen for elevated blood lead levels. Grade: D</p> | | | |
| Risk Assessment | <p>Children younger than age 5 years are at greater risk for elevated blood lead levels and lead toxicity because of increased hand-to-mouth activity, increased lead absorption from the gastrointestinal tract, and the greater vulnerability of the developing central nervous system.</p> <p>Risk factors for increased blood lead levels in children and adults include: minority race/ethnicity; urban residence; low income; low educational attainment; older (pre-1950) housing; recent or ongoing home renovation or remodeling; pica; use of ethnic remedies, certain cosmetics, and exposure to lead-glazed pottery; occupational exposure; and recent immigration. Additional risk factors for pregnant women include alcohol use and smoking.</p> | | | |
| Screening Tests | Venous sampling accurately detects elevated blood lead levels. Screening questionnaires may be of value in identifying children at risk for elevated blood lead levels, but should be tailored for and validated in specific communities for clinical use. | | | |
| Interventions | <p>Treatment options for elevated blood lead levels include residential lead hazard-control efforts (i.e., counseling and education, dust or paint removal, and soil abatement), chelation, and nutritional interventions.</p> <p>Community-based interventions for the prevention of lead exposure are likely to be more effective, and may be more cost-effective, than office-based screening, treatment, and counseling. Relocating children who do not yet have elevated blood lead levels but who live in settings with high lead exposure may be especially helpful.</p> | | | |
| Balance of Benefits and Harms | <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 33%; padding: 5px;"> <p>There is not enough evidence to assess the balance between the potential benefits and harms of routine screening for elevated blood lead levels in children at increased risk.</p> </td> <td style="width: 33%; padding: 5px;"> <p>Given the significant potential harms of treatment and residential lead hazard abatement, and no evidence of treatment benefit, the harms of screening for elevated blood lead levels in children at average risk outweigh the benefits.</p> </td> <td style="width: 33%; padding: 5px;"> <p>Given the significant potential harms of treatment and residential lead hazard abatement, and no evidence of treatment benefit, the harms of screening for elevated blood lead levels in asymptomatic pregnant women outweigh the benefits.</p> </td> </tr> </table> | <p>There is not enough evidence to assess the balance between the potential benefits and harms of routine screening for elevated blood lead levels in children at increased risk.</p> | <p>Given the significant potential harms of treatment and residential lead hazard abatement, and no evidence of treatment benefit, the harms of screening for elevated blood lead levels in children at average risk outweigh the benefits.</p> | <p>Given the significant potential harms of treatment and residential lead hazard abatement, and no evidence of treatment benefit, the harms of screening for elevated blood lead levels in asymptomatic pregnant women outweigh the benefits.</p> |
| <p>There is not enough evidence to assess the balance between the potential benefits and harms of routine screening for elevated blood lead levels in children at increased risk.</p> | <p>Given the significant potential harms of treatment and residential lead hazard abatement, and no evidence of treatment benefit, the harms of screening for elevated blood lead levels in children at average risk outweigh the benefits.</p> | <p>Given the significant potential harms of treatment and residential lead hazard abatement, and no evidence of treatment benefit, the harms of screening for elevated blood lead levels in asymptomatic pregnant women outweigh the benefits.</p> | | |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Child Maltreatment

| | |
|--|--|
| Title | Primary Care Interventions to Prevent Child Maltreatment |
| Population | Children and adolescents aged 0 to 18 years without signs or symptoms of maltreatment |
| Recommendation | No recommendation. Grade: I (Insufficient Evidence) |
| Risk Assessment | <p>There are numerous risk factors associated with child maltreatment, including but not limited to:</p> <ul style="list-style-type: none"> ● Young, single, or nonbiological parents ● Parental lack of understanding of children's needs, child development, or parenting skills ● Poor parent–child relationships/negative interactions ● Parental thoughts or emotions that support maltreatment behaviors ● Family dysfunction or violence ● Parental history of abuse or neglect in family of origin ● Substance abuse within the family ● Social isolation, poverty, or other socioeconomic disadvantages ● Parental stress and distress |
| Interventions | Although the evidence is insufficient to recommend specific preventive interventions, most child maltreatment prevention programs focus on home visitation. Home visitation programs usually comprise a combination of services provided by a nurse or paraprofessional in the family's home on a regularly scheduled basis; most programs are targeted to families with young children and often begin in the prenatal or postnatal period. |
| Balance of Benefits and Harms | The evidence on interventions in primary care to prevent child maltreatment among children without signs or symptoms of maltreatment is insufficient, and the balance of benefits and harms cannot be determined. |
| Other Relevant USPSTF Recommendations | The USPSTF has made recommendations on screening for intimate partner violence and abuse of elderly and vulnerable adults. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Congenital Hypothyroidism

| | |
|-----------------------|---|
| Title | Screening for Congenital Hypothyroidism |
| Population | All newborn infants ¹ |
| Recommendation | Screen for congenital hypothyroidism. Grade: A |

| | |
|---|--|
| Screening Tests | <p>Two methods of screening are used most frequently in the United States:</p> <ul style="list-style-type: none"> ● Primary TSH with backup T4. ● Primary T4 with backup TSH. <p>Screening for congenital hypothyroidism (CH) is mandated in all 50 states and the District of Columbia. Clinicians should become familiar with the tests used in their area and the limitations of the screening strategies employed.</p> |
| Timing of Screening | <p>Infants should be tested between 2 and 4 days of age.</p> <p>Infants discharged from hospitals before 48 hours of life should be tested immediately before discharge. Specimens obtained in the first 24–48 hours of age may be falsely elevated for TSH regardless of the screening method used.</p> |
| Suggestions for Practice | <p>Infants with abnormal screens should receive confirmatory testing and begin appropriate treatment with thyroid hormone replacement within 2 weeks after birth.</p> <p>Children with positive confirmatory testing in whom no permanent cause of CH is found should undergo a 30-day trial of reduced or discontinued thyroid hormone replacement therapy to determine if the hypothyroidism is permanent or transient. This trial of reduced or discontinued therapy should take place at some time after the child reaches 3 years of age.</p> |
| Other Relevant Recommendations from the USPSTF | <p>Additional USPSTF recommendations regarding screening tests for newborns can be accessed at: http://www.uspreventiveservicestaskforce.org/recommendations.htm#pediatric</p> |

¹This recommendation applies to all infants born in the U.S. Premature, very low birth weight and ill infants may benefit from additional screening. These conditions are associated with decreased sensitivity and specificity of screening tests.

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Developmental Dysplasia of the Hip

| | |
|--|--|
| Title | Screening for Developmental Dysplasia of the Hip |
| Population | Infants who do not have obvious hip dislocations or other abnormalities evident without screening |
| Recommendation | No recommendation. Grade: I (Insufficient Evidence) |
| Risk Assessment | Risk factors for developmental dysplasia of the hip include female sex, family history, breech positioning, and in utero postural deformities. However, the majority of cases of developmental dysplasia of the hip have no identifiable risk factors. |
| Screening Tests | Screening tests for developmental dysplasia of the hip have limited accuracy. The most common methods of screening are serial physical examinations of the hip and lower extremities, using the Barlow and Ortolani procedures, and ultrasonography. |
| Interventions | Treatments for developmental dysplasia of the hip include both nonsurgical and surgical options. Nonsurgical treatment with abduction devices is used as early treatment and includes the commonly prescribed Pavlik method. Surgical intervention is used when the dysplasia is severe or diagnosed late, or after an unsuccessful trial of nonsurgical treatment. Avascular necrosis of the hip is the most common and most severe potential harm of both surgical and nonsurgical interventions, and can result in growth arrest of the hip and eventual joint destruction, with significant disability. |
| Balance of Benefits and Harms | The USPSTF was unable to assess the balance of benefits and harms of screening for developmental dysplasia of the hip due to insufficient evidence. There are concerns about the potential harms associated with treatment of infants identified by routine screening. |
| Other Relevant USPSTF Recommendations | The USPSTF has made recommendations on screening for hyperbilirubinemia, phenylketonuria, sickle cell disease, congenital hypothyroidism, and hearing loss in newborns. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Gonococcal Ophthalmia Neonatorum

| | |
|--|--|
| Title | Ocular Prophylaxis for Gonococcal Ophthalmia Neonatorum |
| Population | All newborn infants |
| Recommendation | Provide prophylactic ocular topical medication for the prevention of gonococcal ophthalmia neonatorum. Grade: A |
| Risk Assessment | All newborns should receive prophylaxis. However, some newborns are at increased risk, including those with a maternal history of no prenatal care, sexually transmitted infections, or substance abuse. |
| Preventive Interventions | Preventive medications include 0.5% erythromycin ophthalmic ointment, 1.0% solution of silver nitrate, and 1.0% tetracycline ointment. All are considered equally effective; however, the latter two are no longer available in the United States. |
| Timing of Intervention | Within 24 hours after birth. |
| Other Relevant USPSTF Recommendations | Several recommendations on screening and counseling for infectious diseases and perinatal care can be found at: http://www.uspreventiveservicestaskforce.org . |

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Hearing Loss (Newborns)

| | |
|--|--|
| Title | Universal Screening for Hearing Loss in Newborns |
| Population | All newborn infants |
| Recommendation | Screen for hearing loss in all newborn infants. Grade: B |
| Risk Assessment | <p>The prevalence of hearing loss in newborn infants with specific risk indicators is 10 to 20 times higher than in the general population of newborns.</p> <p>Risk indicators associated with permanent bilateral congenital hearing loss include:</p> <ul style="list-style-type: none"> ● Neonatal intensive care unit admission for 2 or more days. ● Family history of hereditary childhood sensorineural hearing loss. ● Craniofacial abnormalities. ● Certain congenital syndromes and infections. <p>Approximately 50% of newborns with permanent bilateral congenital hearing loss do not have any known risk indicators.</p> |
| Screening Tests | <p>Screening programs should be conducted using a one-step or two-step validated protocol. A frequently-used 2-step screening process involves otoacoustic emissions followed by auditory brain stem response in newborns who fail the first test. Infants with positive screening tests should receive appropriate audiologic evaluation and follow-up after discharge.</p> <p>Procedures for screening and follow-up should be in place for newborns delivered at home, birthing centers, or hospitals without hearing screening facilities.</p> |
| Timing of Screening | All infants should have hearing screening before one month of age. Infants who do not pass the newborn screening should undergo audiologic and medical evaluation before 3 months of age. |
| Treatment | <p>Early intervention services for hearing-impaired infants should meet the individualized needs of the infant and family, including acquisition of communication competence, social skills, emotional well-being, and positive self-esteem.</p> <p>Early intervention comprises evaluation for amplification or sensory devices, surgical and medical evaluation, and communication assessment and therapy. Cochlear implants are usually considered for children with severe-to-profound hearing loss only after inadequate response to hearing aids.</p> |
| Other Relevant USPSTF Recommendations | Additional USPSTF recommendations regarding screening tests for newborns can be accessed at http://www.uspreventiveservicestaskforce.org/recommendations.htm#pediatric . |

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

High Blood Pressure (Children)

| | |
|--|---|
| Title | Screening for Primary Hypertension in Children and Adolescents |
| Population | Children and adolescents without symptoms of hypertension |
| Recommendation | No recommendation. Grade: I (Insufficient Evidence) |
| Risk Assessment | The strongest risk factor for primary hypertension in children is elevated body mass index. Other risk factors include low birthweight, male sex, ethnicity, and a family history of hypertension. |
| Screening Tests | Blood pressure screening with sphygmomanometry in the clinical setting may identify children and adolescents with hypertension with reasonable sensitivity; however, false-positive results may occur with normalization of subsequent blood pressure measurements. |
| Treatment | Stage 1 hypertension in children is treated with lifestyle and pharmacological interventions; medications are not recommended as first-line therapy. |
| Balance of Benefits and Harms | The USPSTF found inadequate evidence on the diagnostic accuracy of screening for primary hypertension. The USPSTF also found inadequate evidence on the effectiveness of treatment and the harms of screening or treatment. Therefore, the USPSTF cannot determine the balance of benefits and harms of screening for hypertension in children and adolescents. |
| Other Relevant USPSTF Recommendations | The USPSTF has made recommendations on screening for lipid disorders in children and adolescents. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Hyperbilirubinemia in Infants

| | |
|--|---|
| Title | Screening of Infants for Hyperbilirubinemia to Prevent Chronic Bilirubin Encephalopathy |
| Population | Healthy newborn infants ≥ 35 weeks' gestational age |
| Recommendation | No recommendation. Grade: I (Insufficient Evidence) |
| Risk Assessment | Risk factors for hyperbilirubinemia include family history of neonatal jaundice, exclusive breastfeeding, bruising, cephalohematoma, ethnicity (Asian, black), maternal age >25 years, male gender, G6PD deficiency, and gestational age <36 weeks. The specific contribution of these risk factors to chronic bilirubin encephalopathy in healthy children is not well understood. |
| Importance | Chronic bilirubin encephalopathy is a rare but devastating condition. Not all children with chronic bilirubin encephalopathy have a history of hyperbilirubinemia. |
| Balance of Benefits and Harms | Evidence about the benefits and harms of screening is lacking. Therefore, the USPSTF could not determine the balance of benefits and harms of screening newborns for hyperbilirubinemia to prevent chronic bilirubin encephalopathy. |
| Considerations for Practice | In deciding whether to screen, clinicians should consider the following: <ul style="list-style-type: none"> ● Potential preventable burden. Bilirubin encephalopathy is a relatively rare disorder. Hyperbilirubinemia alone does not account for the neurologic condition of chronic bilirubin encephalopathy. There is no known screening test that will reliably identify all infants at risk of developing chronic bilirubin encephalopathy. ● Potential harms. Potential harms of screening are unmeasured but may be important. Evidence about the potential harms of phototherapy is lacking. Harms of treatment by exchange transfusion may include apnea, bradycardia, cyanosis, vasospasm, thrombosis, necrotizing enterocolitis, and, rarely, death. ● Current practice. Universal screening is widespread in the United States. |
| Screening Tests | Screening may consist of risk-factor assessment, measurement of bilirubin level either in serum or by transcutaneous estimation, or a combination of methods. |
| Interventions | Phototherapy is commonly used to treat hyperbilirubinemia. Exchange transfusion is used to treat extreme hyperbilirubinemia. |
| Relevant USPSTF Recommendations | USPSTF recommendations on screening newborns for hearing loss, congenital hypothyroidism, hemoglobinopathies, and phenylketonuria (PKU) can be found at http://www.uspreventiveservicestaskforce.org . |

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Illicit and Prescription Drug Use in Children and Adolescents

| | |
|-----------------------|---|
| Title | Primary Care Behavioral Interventions to Reduce Illicit Drug and Nonmedical Pharmaceutical Use in Children and Adolescents |
| Population | Children and adolescents younger than age 18 years who have not already been diagnosed with a substance use disorder |
| Recommendation | No recommendation. Grade: I statement |

| | |
|--|--|
| Behavioral Interventions | While the evidence is insufficient to recommend specific interventions in the primary care setting, those that have been studied include face-to-face counseling, videos, print materials, and interactive computer-based tools. Studies on these interventions were limited and findings on whether interventions significantly improved health outcomes were inconsistent. |
| Balance of Benefits and Harms | The evidence regarding primary care–based behavioral interventions to prevent or reduce illicit drug and nonmedical pharmaceutical use in children and adolescents is insufficient, and the balance of benefits and harms cannot be determined. |
| Other Relevant USPSTF Recommendations | The USPSTF has made recommendations on screening for and interventions to decrease the unhealthy use of other substances, including alcohol and tobacco. These recommendations are available at www.uspreventiveservicestaskforce.org . |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Iron Deficiency Anemia (Screening)

| | | |
|-----------------------|--|-----------------------------|
| Title | Part I: Screening for Iron Deficiency Anemia in Children and Pregnant Women | |
| Population | Asymptomatic children ages 6 to 12 months | Asymptomatic pregnant women |
| Recommendation | Screen for iron deficiency anemia. Grade: B | |

| | | |
|--|---|---|
| Risk Assessment | Individuals considered to be at high risk for iron deficiency include adult women, recent immigrants, and, among adolescent females, fad dieters, as well as those who are obese. Premature and low birth weight infants are also at increased risk for iron deficiency. | |
| Screening Tests | Serum hemoglobin or hematocrit is the primary screening test for identifying anemia. Hemoglobin is sensitive for iron deficiency anemia; however, it is not sensitive for iron deficiency because mild deficiency states may not affect hemoglobin levels. Potential harms of screening include false-positive results, anxiety, and cost. | |
| Interventions | Iron deficiency anemia is usually treated with oral iron preparations. The likelihood that iron deficiency anemia identified by screening will respond to treatment is unclear, because many families do not adhere to treatment and because the rate of spontaneous resolution is high. | |
| Balance of Benefits and Harms | The USPSTF was unable to determine the balance between the benefits and harms of routine screening for iron deficiency anemia in asymptomatic children ages 6 to 12 months. | The benefits of routine screening for iron deficiency anemia in asymptomatic pregnant women outweigh the potential harms. |
| Other Relevant USPSTF Recommendations | The USPSTF has also made recommendations on screening for blood lead levels in children and pregnant women. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . | |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Iron Deficiency Anemia (Supplementation)

| Title | Part II: Iron Supplementation for Children and Pregnant Women | | |
|-----------------------|--|--|--|
| Population | Asymptomatic children ages 6 to 12 months who are at increased risk for iron deficiency anemia | Asymptomatic children ages 6 to 12 months who are at average risk for iron deficiency anemia | Pregnant women who are not anemic |
| Recommendation | Provide routine iron supplementation. Grade: B | No recommendation. Grade: I (Insufficient Evidence) | No recommendation. Grade: I (Insufficient Evidence) |

| | | | |
|--|---|---|---|
| Risk Assessment | A validated risk assessment tool to guide primary care physicians in identifying individuals who would benefit from iron supplementation has not been developed. | | |
| Preventive Medication | Iron supplementation, such as iron-fortified formula or iron supplements, may improve neurodevelopmental outcomes in children at increased risk for iron deficiency anemia. There is poor evidence that it improves neurodevelopmental or health outcomes in other populations. Oral iron supplementation increases the risk for unintentional overdose and gastrointestinal symptoms. Given appropriate protection against overdose, these harms are small. | | |
| Balance of Benefits and Harms | The moderate benefits of iron supplementation in asymptomatic children ages 6 to 12 months who are at increased risk for iron deficiency anemia outweigh the potential harms. | The USPSTF was unable to determine the balance between the benefits and harms of iron supplementation in children ages 6 to 12 months who are at average risk for iron deficiency anemia. | The USPSTF was unable to determine the balance between the benefits and harms of iron supplementation in non-anemic pregnant women. |
| Other Relevant USPSTF Recommendations | The USPSTF has also made recommendations on folic acid supplementation in women planning or capable of pregnancy and vitamin D supplementation to prevent cancer and fractures. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . | | |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Lipid Disorders in Children

| | |
|--|--|
| Title | Screening for Lipid Disorders in Children |
| Population | Asymptomatic infants, children, adolescents, and young adults (age 20 years or younger) |
| Recommendation | No recommendation. Grade: I (Insufficient Evidence) |
| Risk Assessment | Risk factors for dyslipidemia include overweight, diabetes, and a family history of common familial dyslipidemias (e.g., familial hypercholesterolemia). |
| Screening Tests | Serum lipid (total cholesterol, high-density and low-density lipoprotein cholesterol) levels are accurate screening tests for childhood dyslipidemia, although many children with multifactorial types of dyslipidemia will have normal lipid levels in adulthood. The use of family history as a screening tool for dyslipidemia has variable accuracy, largely because definitions of a positive family history and lipid threshold values vary substantially. |
| Interventions | The effectiveness of treatment interventions (diet, exercise, lipid-lowering agents) in improving health outcomes in children with dyslipidemia (including multifactorial dyslipidemia) remains a critical research gap. Potential harms of screening may include labeling of children whose dyslipidemia would not persist into adulthood or cause health problems. Adverse effects from lipid-lowering medications and low-fat diets, including potential long-term harms, have been inadequately evaluated in children. |
| Balance of Benefits and Harms | The USPSTF was unable to determine the balance between the potential benefits and harms of routinely screening children and adolescents for dyslipidemia. |
| Other Relevant USPSTF Recommendations | The USPSTF has made recommendations on high blood pressure and obesity in children and adolescents. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Major Depressive Disorder in Children and Adolescents

| | | |
|-----------------------|--|-----------------------|
| Title | Screening and Treatment for Major Depressive Disorder in Children and Adolescents | |
| Population | Adolescents (12-18 years) | Children (7-11 years) |
| Recommendation | Screen when systems for diagnosis, treatment, and followup are in place. Grade: B | |
| | No Recommendation Grade: I (Insufficient Evidence) | |

| | | |
|------------------------|---|--|
| Risk Assessment | Risk factors for major depressive disorder (MDD) include parental depression, having comorbid mental health or chronic medical conditions, and having experienced a major negative life event. | |
| Screening Tests | <p>The following screening tests have been shown to do well in teens in primary care settings:</p> <ul style="list-style-type: none"> • Patient Health Questionnaire for Adolescents (PHQ-A). • Beck Depression Inventory-Primary Care Version (BDI-PC). | Screening instruments perform less well in younger children. |
| Treatments | Among pharmacotherapies fluoxetine, a selective serotonin reuptake inhibitor (SSRI), has been found efficacious. However, because of risk of suicidality, SSRIs should be considered only if clinical monitoring is possible. Various modes of psychotherapy, and pharmacotherapy combined with psychotherapy, have been found efficacious. | Evidence on the balance of benefits and harms of treatment of younger children is insufficient for a recommendation. |

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Obesity in Children and Adolescents

| | |
|-----------------------|---|
| Title | Screening for Obesity in Children and Adolescents |
| Population | Children and adolescents 6 to 18 years of age |
| Recommendation | Screen children aged 6 years and older for obesity. Offer or refer for intensive counseling and behavioral interventions. Grade: B |

| | |
|--|--|
| Screening Tests | BMI is calculated from the weight in kilograms divided by the square of the height in meters. Height and weight, from which BMI is calculated, are routinely measured during health maintenance visits. BMI percentile can be plotted on a chart or obtained from online calculators. Overweight age- and gender-specific BMI at ≥85th to 94th percentile Obesity age- and gender-specific BMI at ≥95th percentile |
| Timing of Screening | No evidence was found on appropriate screening intervals. |
| Interventions | Refer patients to comprehensive moderate- to high-intensity programs that include dietary, physical activity, and behavioral counseling components. |
| Balance of Benefits and Harms | Moderate- to high-intensity programs were found to yield modest weight changes. Limited evidence suggests that these improvements can be sustained over the year after treatment. Harms of screening were judged to be minimal. |
| Other Relevant USPSTF Recommendations | Recommendations on other pediatric and behavioral counseling topics can be found at http://www.uspreventiveservicestaskforce.org . |

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Phenylketonuria (PKU)

| | |
|--|---|
| Title | Screening for Phenylketonuria |
| Population | All newborn infants |
| Recommendation | Screen for Phenylketonuria (PKU). Grade: A |
| Screening Tests | Screening for PKU is mandated in all 50 states. Methods of screening vary. Three main methods are used to screen for PKU in the United States: <ol style="list-style-type: none"> 1. Guthrie Bacterial Inhibition Assay (BIA) 2. Automated fluorometric assay 3. Tandem mass spectrometry |
| Timing of Screening | Infants who are tested within the first 24 hours after birth should receive a repeat screening test by 2 weeks of age. Optimal timing of screening for premature infants and infants with illnesses is at or near 7 days of age, but in all cases before discharge from the newborn nursery. |
| Treatment | It is essential that phenylalanine restrictions be instituted shortly after birth to prevent the neurodevelopmental effects of PKU. |
| Other Relevant USPSTF Recommendations | Additional USPSTF recommendations regarding screening tests for newborns can be accessed at: http://www.uspreventiveservicestaskforce.org/recommendations.htm#pediatric |

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Sickle Cell Disease

| | |
|--|---|
| Title | Screening for Sickle Cell Disease in Newborns |
| Population | All newborn infants |
| Recommendation | Screen for sickle cell disease. Grade: A |
| Screening Tests | <p>Screening for sickle cell disease in newborns is mandated in all 50 states and the District of Columbia. In most states, one of these tests is used for the initial screening:</p> <ul style="list-style-type: none"> • Thin-layer isoelectric focusing (IEF). • High performance liquid chromatography (HPLC). <p>Both IEF and HPLC have extremely high sensitivity and specificity for sickle cell anemia.</p> |
| Timing of Screening | <p>All newborns should undergo screening regardless of birth setting. Birth attendants should make arrangements for samples to be obtained. The first clinician to see the infant at an office visit should verify screening results. Confirmatory testing should occur no later than 2 months of age.</p> |
| Treatment | <p>Infants with sickle cell anemia should receive:</p> <ul style="list-style-type: none"> • Prophylactic penicillin starting by age 2 months. • Pneumococcal immunizations at recommended intervals. |
| Other Relevant USPSTF Recommendations | <p>Additional USPSTF recommendations regarding screening tests for newborns can be accessed at http://www.uspreventiveservicestaskforce.org/recommendations.htm#vision.</p> |

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.

Speech and Language Delay

| | |
|-----------------------|--|
| Title | Screening for Speech and Language Delay in Preschool Children |
| Population | Children ages 5 years and younger who have not already been identified as at increased risk for speech and language delays |
| Recommendation | No recommendation. Grade: I (Insufficient Evidence) |

| | |
|--|--|
| Risk Assessment | The most consistently reported risk factors include a family history of speech and language delay, male sex, and perinatal factors, such as prematurity and low birth-weight. Other risk factors reported less consistently include levels of parental education, specific childhood illnesses, birth order, and larger family size. |
| Screening Tests | There is insufficient evidence that brief, formal screening instruments that are suitable for use in primary care for assessing speech and language development can accurately identify children who would benefit from further evaluation and intervention. |
| Balance of Benefits and Harms | The USPSTF could not determine the balance of benefits and harms of using brief, formal screening instruments to screen for speech and language delay in the primary care setting. |
| Other Relevant USPSTF Recommendations | The USPSTF has also made recommendations on screening for hearing loss in newborns and vision impairment in children ages 1 to 5 years. These recommendations are available at http://www.uspreventiveservicestaskforce.org/ . |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Tobacco Use in Children and Adolescents


| | |
|--|--|
| Title | Primary Care Interventions to Prevent Tobacco Use in Children and Adolescents |
| Population | School-aged children and adolescents |
| Recommendation | Provide interventions to prevent initiation of tobacco use. Grade: B |
| Risk Assessment | The strongest factors associated with smoking initiation in children and adolescents are parental smoking and parental nicotine dependence. Other factors include low levels of parental monitoring, easy access to cigarettes, perception that peers smoke, and exposure to tobacco promotions. |
| Behavioral Counseling Interventions | Behavioral counseling interventions, such as face-to-face or phone interaction with a health care provider, print materials, and computer applications, can reduce the risk for smoking initiation in school-aged children and adolescents. The type and intensity of effective behavioral interventions substantially varies. |
| Balance of Benefits and Harms | There is a moderate net benefit to providing primary care interventions to prevent tobacco use in school-aged children and adolescents. |
| Other Relevant USPSTF Recommendations | The USPSTF has made recommendations on counseling and interventions to prevent tobacco use and tobacco-caused disease in adults and pregnant women. These recommendations are available at www.uspreventiveservicestaskforce.org/ . |

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org/>.

Visual Impairment in Children Ages 1 to 5

| Screening for Visual Impairment in Children Ages 1 to 5 | |
|---|--|
| Title | Screening for Visual Impairment in Children Ages 1 to 5 |
| Population | Children ages 3 to 5 years |
| Recommendation | <p style="text-align: center;">Children younger than 3 years of age</p> <p style="text-align: center;">No recommendation.</p> <p style="text-align: center;">Grade: I (Insufficient Evidence)</p> |
| Screening Tests | <p>Various screening tests are used in primary care to identify visual impairment in children, including:</p> <ul style="list-style-type: none"> ● Visual acuity test ● Stereoacuity test ● Cover-uncover test ● Hirschberg light reflex test ● Autorefraction ● Photoscreening |
| Timing of Screening | No evidence was found regarding appropriate screening intervals. |
| Interventions | Primary treatment for amblyopia includes the use of corrective lenses, patching, or atropine therapy of the non-affected eye. Treatment may also consist of a combination of interventions. |
| Balance of Benefits and Harms | <p>There is adequate evidence that early treatment of amblyopia in children ages 3 to 5 years leads to improved visual outcomes. There is limited evidence on harms of screening, including psychosocial effects, in children ages 3 years and older.</p> <p>There is inadequate evidence that early treatment of amblyopia in children younger than 3 years of age leads to improved visual outcomes.</p> |
| Suggestions for Practice Regarding the I Statement | <p>In deciding whether to refer children younger than 3 years of age for screening, clinicians should consider:</p> <ul style="list-style-type: none"> ● <i>Potential preventable burden</i>: screening later in the preschool years seems to be as effective as screening earlier ● <i>Costs</i>: initial high costs associated with autorefractors and photoscreeners ● <i>Current practice</i>: typical vision screening includes assessment of visual acuity, strabismus, and stereoacuity; children with positive findings should be referred for a comprehensive ophthalmologist exam |

For a summary of the evidence systematically reviewed in making these recommendations, the full recommendation statement, and supporting documents, please go to <http://www.uspreventiveservicestaskforce.org>.



**Immunizations
for Adults and
Children**

Immunizations for Adults and Children

The USPSTF recognizes the importance of immunizations in primary disease prevention. However, the USPSTF does not wish to duplicate the significant investment of resources made by others to review new evidence on immunizations in a timely fashion and make recommendations.

The Centers for Disease Control and Prevention's (CDC's) Advisory Committee on Immunization Practices (ACIP) publishes recommendations on immunizations for children and adults. The methods used by the ACIP to review evidence on immunizations may differ from the methods used by the USPSTF.

For the ACIP's current recommendations on immunizations, please refer to the National Immunization Program Web site at www.cdc.gov/vaccines/recs/schedules/default.htm.

The background of the slide is a light gray color with several large, overlapping, curved bands in a slightly darker shade of gray. These bands sweep across the frame from the left side towards the right, creating a sense of motion and depth. The text is centered within the space created by these curves.

Topics in Progress

Topics in Progress

Each USPSTF recommendation goes through several stages of development. The review process takes into account input from the medical and research community, stakeholders, and the general public.

The length of time for the entire recommendation process varies depending on the amount and type of available evidence and the time required for compilation of data into a draft recommendation, public comment periods, consideration of comments, and in-depth review and discussions among USPSTF members.

The following topics are in review and are likely to be issued as drafts for public comment during 2014:

- Aspirin for the Prevention of Cardiovascular Disease
- Aspirin or NSAIDS for the Prevention of Colorectal Cancer, Preventive Medications
- Aspirin Use to Prevent Preeclampsia, Preventive Medications
- Chlamydial Infection, Screening
- Diabetes Mellitus, Screening
- Gonorrhea, Screening
- Healthy Diet and Physical Activity for Cardiovascular Disease Prevention in Adults at Risk, Counseling
- High Blood Pressure in Adults, Screening
- Iron Deficiency Anemia in Childhood and Pregnancy, Screening
- Major Depressive Disorder in Children and Adolescents, Screening
- Sexually Transmitted Infections, Counseling
- Thyroid Disease, Screening
- Vitamin D Deficiency, Screening

Recommendations on the following topics were published during the production of the *2014 Guide to Clinical Preventive Services* or are in review and are likely to be published as final recommendations during 2014:

- Abdominal Aortic Aneurysm, Screening
- Carotid Artery Stenosis, Screening
- Dental Caries Prevention in Children From Birth to Age 5 Years, Screening and Preventive Medications

- Hepatitis B Virus Infection in Nonpregnant Adolescents and Adults, Screening
- Suicide Risk, Screening

The following topics are in earlier stages of review and are likely to be issued as drafts or published as final recommendations sometime after 2014:

- Autism Spectrum Disorder in Young Children, Screening
- Breast Cancer, Screening
- Chronic Obstructive Pulmonary Disease, Screening
- Colorectal Cancer, Screening
- Depression in Adults and in Pregnant and Postpartum Women, Screening
- Impaired Visual Acuity in Older Adults, Screening
- Lipid Disorders in Adults, Screening
- Lipid Disorders in Children, Screening
- Skin Cancer, Screening
- Speech and Language Delay and Disorders in Children Age 5 Years or Younger, Screening
- Syphilis Infection, Screening
- Tobacco Cessation in Adults, Including Pregnant Women, Counseling

The background of the page is a light gray color. On the left side, there are several thick, curved, overlapping lines in a slightly darker shade of gray, creating a sense of motion or a stylized wave pattern. The lines curve from the top left towards the bottom right.

Appendixes and Index

Appendix A

How the U.S. Preventive Services Task Force Grades Its Recommendations

The Task Force assigns each of its recommendations a letter grade (A, B, C, or D) or issues an I statement, based on the certainty of the evidence and on the balance of benefits and harms of the preventive service as displayed in the recommendation grid below. The USPSTF changed its grade definitions based on a change in methods in May 2007.

U.S. Preventive Services Task Force Recommendation Grid: Letter Grade of Recommendation or Statement of Insufficient Evidence Assessing Certainty and Magnitude of Net Benefit

| Certainty of Net Benefit | Magnitude of Net Benefit | | | |
|--------------------------|--------------------------|----------|-------|---------------|
| | Substantial | Moderate | Small | Zero/negative |
| High | A | B | C | D |
| Moderate | B | B | C | D |
| Low | Insufficient | | | |

Grade Definitions

What the Grades Mean and Suggestions for Practice

The USPSTF updated its definitions of the grades it assigns to recommendations and now includes “suggestions for practice” associated with each grade. The USPSTF has also defined levels of certainty regarding net benefit.

| Grade | Definition | Suggestions for Practice |
|--------------------|--|---|
| A | The USPSTF recommends the service. There is high certainty that the net benefit is substantial. | Offer or provide this service. |
| B | The USPSTF recommends the service. There is high certainty that the net benefit is moderate or there is moderate certainty that the net benefit is moderate to substantial. | Offer or provide this service. |
| C | The USPSTF recommends selectively offering or providing this service to individual patients based on professional judgment and patient preferences. There is at least moderate certainty that the net benefit is small. | Offer or provide this service for selected patients depending on individual circumstances. |
| D | The USPSTF recommends against the service. There is moderate or high certainty that the service has no net benefit or that the harms outweigh the benefits. | Discourage the use of this service. |
| I Statement | The USPSTF concludes that the current evidence is insufficient to assess the balance of benefits and harms of the service. Evidence is lacking, of poor quality, or conflicting, and the balance of benefits and harms cannot be determined. | Read the clinical considerations section of USPSTF Recommendation Statement. If the service is offered, patients should understand the uncertainty about the balance of benefits and harms. |

Levels of Certainty Regarding Net Benefit

| Level of Certainty* | Description |
|---------------------|--|
| High | <p>The available evidence usually includes consistent results from well-designed, well-conducted studies in representative primary care populations. These studies assess the effects of the preventive service on health outcomes. This conclusion is therefore unlikely to be strongly affected by the results of future studies.</p> |
| Moderate | <p>The available evidence is sufficient to determine the effects of the preventive service on health outcomes, but confidence in the estimate is constrained by such factors as:</p> <ul style="list-style-type: none"> ● The number, size, or quality of individual studies. ● Inconsistency of findings across individual studies. ● Limited generalizability of findings to routine primary care practice. ● Lack of coherence in the chain of evidence. <p>As more information becomes available, the magnitude or direction of the observed effect could change, and this change may be large enough to alter the conclusion.</p> |
| Low | <p>The available evidence is insufficient to assess effects on health outcomes. Evidence is insufficient because of:</p> <ul style="list-style-type: none"> ● The limited number or size of studies. ● Important flaws in study design or methods. ● Inconsistency of findings across individual studies. ● Gaps in the chain of evidence. ● Findings not generalizable to routine primary care practice. ● Lack of information on important health outcomes. <p>More information may allow estimation of effects on health outcomes.</p> |

* The USPSTF defines certainty as “likelihood that the USPSTF assessment of the net benefit of a preventive service is correct.” The net benefit is defined as benefit minus harm of the preventive service as implemented in a general, primary care population. The USPSTF assigns a certainty level based on the nature of the overall evidence available to assess the net benefit of a preventive service.

Grade Definitions Prior to May 2007

The definitions below (of USPSTF grades and quality of evidence ratings) were in use prior to the update and apply to recommendations voted on by the USPSTF prior to May 2007.

A — Strongly Recommended: The USPSTF strongly recommends that clinicians provide [the service] to eligible patients. The USPSTF found good evidence that [the service] improves important health outcomes and concludes that benefits substantially outweigh harms.

B — Recommended: The USPSTF recommends that clinicians provide [the service] to eligible patients. The USPSTF found at least fair evidence that [the service] improves important health outcomes and concludes that benefits outweigh harms.

C — No Recommendation: The USPSTF makes no recommendation for or against routine provision of [the service]. The USPSTF found at least fair evidence that [the service] can improve health outcomes but concludes that the balance of benefits and harms is too close to justify a general recommendation.

D — Not Recommended: The USPSTF recommends against routinely providing [the service] to asymptomatic patients. The USPSTF found at least fair evidence that [the service] is ineffective or that harms outweigh benefits.

I — Insufficient Evidence to Make a Recommendation: The USPSTF concludes that the evidence is insufficient to recommend for or against routinely providing [the service]. Evidence that [the service] is effective is lacking, of poor quality, or conflicting and the balance of benefits and harms cannot be determined.

Quality of Evidence Prior to May 2007

The USPSTF grades the quality of the overall evidence for a service on a 3-point scale (good, fair, poor):

Good: Evidence includes consistent results from well-designed, well-conducted studies in representative populations that directly assess effects on health outcomes.

Fair: Evidence is sufficient to determine effects on health outcomes, but the strength of the evidence is limited by the number, quality, or consistency of the individual studies, generalizability to routine practice, or indirect nature of the evidence on health outcomes.

Poor: Evidence is insufficient to assess the effects on health outcomes because of limited number or power of studies, important flaws in their design or conduct, gaps in the chain of evidence, or lack of information on important health outcomes.

Appendix B

Members of the U.S. Preventive Services Task Force 2004-2014

**Janet D. Allan, Ph.D., R.N., C.S.,
F.A.A.N.**

School of Nursing
University of Maryland, Baltimore
Baltimore, MD

Linda Ciofu Baumann, Ph.D., R.N.

School of Nursing and School of
Medicine & Public Health
University of Wisconsin
Madison, WI

Alfred O. Berg, M.D., M.P.H.

University of Washington
Seattle, WA

**Kirsten Bibbins-Domingo, Ph.D.,
M.D.**

San Francisco General Hospital
University of California,
San Francisco, CA

Adelita Gonzales Cantu, R.N., Ph.D.

University of Texas Health Science
Center
San Antonio, TX

Ned Calonge, M.D., M.P.H.

Colorado Department of Public Health
and Environment
Denver, CO

Susan J. Curry, Ph.D.

College of Public Health
University of Iowa
Iowa City, IA

Karina W. Davidson, Ph.D., M.A.Sc.

Center for Behavioral Cardiovascular
Health
Columbia University Medical Center
New York, NY

Thomas G. DeWitt, M.D.

Department of Pediatrics
Children's Hospital Medical Center
Cincinnati, OH

Allen J. Dietrich, M.D.

Dartmouth Medical School
Hanover, NH

Mark Ebell, M.D., M.S.

The University of Georgia
Athens, GA

Glenn Flores, M.D.

University of Texas
Southwestern Medical Center and
Children's Medical Center of Dallas
Dallas, TX

Paul S. Frame, M.D.

Tri-County Family Medicine
Cohocton, NY

Francisco A.R. García, M.D., M.P.H.

Pima County Department of Health
Tucson, AZ

Matthew W. Gillman, M.D., S.M.

Obesity Prevention Program
Harvard Medical School and Harvard
Pilgrim Health Care Institute
Boston, MA

Leon Gordis, M.D., Dr.P.H.

Johns Hopkins Bloomberg School of
Public Health
Baltimore, MD

Kimberly D. Gregory, M.D., M.P.H.

Cedars-Sinai Medical Center
Los Angeles, CA

David Grossman, M.D., M.P.H.

Center for Health Studies, Group
Health Cooperative
University of Washington
Seattle, WA

Russell Harris, M.D., M.P.H.

University of North Carolina School
of Medicine
Chapel Hill, NC

Jessica Herzstein, M.D., M.P.H.

Air Products
Allentown, PA

George Isham, M.D., M.S.

HealthPartners
Minneapolis, MN

Mark S. Johnson, M.D., M.P.H.

New Jersey Medical School
University of Medicine and Dentistry of
New Jersey
Newark, NJ

Alex R. Kemper, M.D., M.P.H., M.S.

Pediatrics
Duke University Medical School
Durham, NC
Division of Children's Primary Care
Duke University
Durham, NC

Kenneth Kizer, M.D., M.P.H.

National Quality Forum
Washington, DC

Jonathan D. Klein, M.D., M.P.H.

University of Rochester
Rochester, NY

**Ann E. Kurth, Ph.D., R.N., M.S.N.,
M.P.H.**

College of Nursing and the Department
of Population Health
Global Institute of Public Health
New York University
New York, NY

Michael L. LeFevre, M.D., M.S.P.H.

University of Missouri School of
Medicine
Columbia, MO

Rosanne Leipzig, M.D., Ph.D.

Mount Sinai School of Medicine
New York, NY

**Carol Loveland-Cherry, Ph.D., R.N.,
F.A.A.N.**

School of Nursing
University of Michigan
Ann Arbor, MI

Lucy N. Marion, Ph.D., R.N.

School of Nursing
Medical College of Georgia
Augusta, GA

Joy Melnikow, M.D., M.P.H.

University of California Davis
Sacramento, CA

**Bernadette Melnyk, Ph.D., R.N.,
C.P.N.P./N.P.P.**

College of Nursing & Healthcare
Innovation
Arizona State University
Phoenix, AZ

Virginia A. Moyer, M.D., M.P.H.

University of Texas Health Science
Center
Houston, TX

**Wanda Nicholson, M.D., M.P.H.,
M.B.A.**

Johns Hopkins School of Medicine and
Bloomberg School of Public Health
Baltimore, MD

Judith K. Ockene, Ph.D., M.Ed.

University of Massachusetts Medical
School
Worcester, MA

Douglas K. Owens, M.D., M.S.
VA Palo Alto Health Care System
Freeman Spogli Institute for
International Studies
Stanford University
Stanford, CA

Diana B. Petitti, M.D., M.P.H.
Fulton School of Engineering
Arizona State University
Tempe, AZ

William R. Phillips, M.D., M.P.H.
Family Medicine and Health Services
University of Washington
Seattle, WA

Maureen G. Phipps, M.D., M.P.H.
Department of Obstetrics and
Gynecology
Warren Alpert Medical School
Brown University, Providence, RI

Michael P. Pignone, M.D., M.P.H.
Division of General Internal Medicine
University of North Carolina
Chapel Hill, NC

Carolina Reyes, M.D.
University of Southern California,
Los Angeles County/USC Medical
Center
Los Angeles, CA

George F. Sawaya, M.D.
University of California, San Francisco
San Francisco, CA

J. Sanford (Sandy) Schwartz, M.D.
University of Pennsylvania School of
Medicine and Wharton School
Philadelphia, PA

Albert L. Siu, M.D.
Mount Sinai Medical Center
New York, NY

Steven M. Teutsch, M.D., M.P.H.
Merck and Company, Inc.
West Point, PA

Timothy Wilt, M.D., M.P.H.
Minneapolis VA Medical Center
University of Minnesota
Minneapolis, MN

**Barbara P. Yawn, M.D., M.S.P.H.,
M.Sc.**
Olmstead Medical Center
Rochester, MN

Appendix C

Acknowledgements

AHRQ Staff Supporting the USPSTF

Aileen Buckler, M.D., M.P.H.
Joel Boches
Joya Chowdhury, M.P.H.
Robert Cosby, Ph.D.
Jennifer Crowell, M.D., M.P.H.
Sandra K. Cummings
Farah Englert
Tina Fan, M.D., M.P.H.
Saeed Fatemi
Janice Genevro, Ph.D., M.S.W.
Margi Grady
Alison Hunt
William Hyde, M.L.S.
Kristie Kiser
Karen Lee, M.D., M.P.H.
Biff LeVee
Iris Mabry, M.D., M.P.H.
Corey Mackison, M.S.A.
Andrew Marshall
Robert McNellis, M.P.H., P.A.
David Meyers, M.D.
Tess Miller, Dr.P.H.
Emily Moser
Quyen Ngo-Metzger, M.D., M.P.H.
Lisa Nicolella
Linwood Norman, M.S.
Janine Payne, M.P.H.
Kathryn Ramage
Richard Ricciardi, Ph.D., N.P., F.A.A.N.P.
Randie Siegel, M.S.
Rachel Weinstein
Tracy Wolff, M.D., M.P.H.

Evidence-based Practice Centers Supporting the USPSTF 2014

Kaiser Permanente Research Affiliates Evidence-based Practice Center

Director, Evelyn P. Whitlock, M.D., M.P.H.

The Kaiser Permanente Research Affiliates EPC involves over 75 researchers from three integrated health care delivery systems: Kaiser Permanente, Group Health Research Institute, and HealthPartners Institute for Education and Research. The KPRA EPC is headquartered in Portland, Oregon, at the Kaiser Permanente Center for Health Research. To learn more about the KPRA EPC, please visit:

<http://www.kpchr.org/research/public/resEPC.htm>

Pacific Northwest Evidence-based Practice Center

Director, Roger Chou, M.D.

The Pacific Northwest EPC is housed in the Department of Medical Informatics & Clinical Epidemiology, at the Oregon Health and Science University School of Medicine, with partners at the University of Washington CHASE Alliance in Seattle, and Spectrum Research, Inc., in Tacoma, Washington. To learn more about the Pacific Northwest EPC, please visit:

<http://www.ohsu.edu/xd/research/centers-institutes/evidence-based-practice-center/>

RTI-UNC Evidence-based Practice Center

Director, Meena Viswanathan, Ph.D.

Co-Director, Dan Jonas, M.D., M.P.H.

RTI International, in collaboration with the five health professions schools and the Cecil G. Sheps Center for Health Services Research at the University of North Carolina at Chapel Hill, operates the RTI-UNC Evidence-based Practice Center. To learn more about the RTI-UNC EPC, please visit:

<http://www.rti.org/page.cfm?objectid=4FA3323C-51CA-4DFE-A3E0246F90D5FA4B>.

Vanderbilt Evidence-based Practice Center

Director, Melissa McPheeters, Ph.D., M.P.H.

Established in 2007, the Vanderbilt EPC is located in the Institute for Medicine and Public Health at Vanderbilt University. To learn more about the Vanderbilt EPC, please visit:

<http://medicineandpublichealth.vanderbilt.edu/epc/>.

Liaisons to the USPSTF

Primary care partners include:

- American Academy of Family Physicians (AAFP)
- American Academy of Nurse Practitioners (AANP)
- American Academy of Pediatrics (AAP)
- American Academy of Physician Assistants (AAPA)
- American Congress of Obstetricians and Gynecologists (ACOG)
- American College of Physicians (ACP)
- American College of Preventive Medicine (ACPM)
- American Osteopathic Association (AOA)
- National Association of Pediatric Nurse Practitioners (NAPNAP)

Policy, population, and quality improvement partners include:

- America's Health Insurance Plans (AHIP)
- American Medical Association
- National Committee for Quality Assurance (NCQA)
- AARP
- Canadian Task Force on Preventive Health Care (CTFPHC)
- Community Preventive Services Task Force (CPSTF)

Federal partners include:

- Centers for Disease Control and Prevention (CDC)
- Centers for Medicare & Medicaid Services (CMS)
- Department of Defense/Military Health System (DoD/MHS)
- U.S. Food and Drug Administration (FDA)
- Health Resources and Services Administration (HRSA)
- Indian Health Service (IHS)
- National Institutes of Health (NIH)
- Veteran's Health Administration (VHA)
- Substance Abuse & Mental Health Services Administration (SAMHSA)
- Office of the Assistant Secretary for Planning and Evaluation (ASPE)
- Office of Disease Prevention and Health Promotion (ODPHP)
- Office of the Surgeon General

Appendix D

About the U.S. Preventive Services Task Force

Overview

Created in 1984, the U.S. Preventive Services Task Force (USPSTF) is an independent group of national experts in prevention and evidence-based medicine that works to improve the health of all Americans by making evidence-based recommendations about clinical preventive services such as:

- Screenings
- Counseling services
- Preventive medications

The Task Force is made up of 16 volunteer members who serve 4-year terms. Members come from the fields of preventive medicine and primary care, including internal medicine, family medicine, pediatrics, behavioral health, obstetrics and gynecology, and nursing. The Task Force is led by a chair and two vice-chairs. Members are appointed by the Director of the Agency for Healthcare Research and Quality (AHRQ). Members must have no substantial conflicts of interest that could impair the integrity of the work of the Task Force. A list of current USPSTF members, including their biographical information, can be found on the USPSTF Web site (www.USPreventiveServicesTaskForce.org).

Since 1998, through acts of the U.S. Congress, AHRQ has been authorized to convene the Task Force and to provide ongoing scientific, administrative, and dissemination support to the Task Force.

Recommendations

The Task Force makes recommendations to help primary care clinicians and patients decide together whether a preventive service is right for a patient's needs. Its recommendations apply to people who have no signs or symptoms of the specific disease or condition to which a recommendation applies and are for services prescribed, ordered, or delivered in the primary care setting.

Task Force recommendations are based on a rigorous review of existing peer-reviewed evidence. The Task Force assesses the effectiveness of a clinical preventive service by evaluating and balancing the potential benefits and harms of the service. The potential benefits include early identification of disease leading to improvement in health. The potential harms can include adverse effects of the service itself or inaccurate test results that may lead to additional testing, additional risks, or unneeded treatment.

The Task Force does not explicitly consider costs in its assessment of the effectiveness of a service. The Task Force assigns each recommendation a letter grade (A, B, C, or D grade or an I statement) based on the strength of the evidence and on the balance of benefits and harms of the preventive service. More information on USPSTF recommendation grades and a list of all current USPSTF recommendations can be found on the USPSTF Web site.

The Recommendation Making Process

The USPSTF is committed to making its work as transparent as possible. As part of this commitment, the Task Force provides opportunities for the public to provide input during each phase of the recommendation process.

The phases of the topic development process are described below and illustrated in “Steps the USPSTF Takes to Make a Recommendation” at the end of this appendix.

Topic Nomination

The USPSTF considers a broad range of clinical preventive services for its recommendations, focusing on screenings, counseling, and preventive medications. Anyone can nominate a topic for consideration by the Task Force.

Research Plan Development

Once the USPSTF selects a topic for review, it works with an Evidence-based Practice Center (EPC) to develop a draft research plan, which guides the review process and includes key questions and target populations. A draft research plan is posted for public comment, and feedback is incorporated into a final research plan.

Evidence Report and Recommendation Statement Development

Using the final research plan as a guide, the EPC researchers independently gather, review, and analyze evidence on the topic and summarize their findings in a systematic evidence report. The evidence report is sent to subject matter experts for peer review before it is shared with the Task Force.

Then, the entire Task Force discusses and deliberates the evidence, weighs the benefits and harms, and uses the information to determine the effectiveness of a service. The Task Force revises and finalizes a draft recommendation statement based on this discussion.

The draft evidence report and draft recommendation statement are typically posted together on the Task Force Web site for a period of 4 weeks. During the comment period, any member of the public may submit comments on either or both of the documents.

Upon receiving all comments from experts, partners, and the public, the EPC researchers revise the draft evidence report. Once final, the EPC begins to prepare a summary of the evidence report for submission to a peer-reviewed journal.

Final Recommendation Statement

Then, Task Force members review all the comments received and use them to inform the development of the final recommendation statement. The recommendation statement is sent to all Task Force members for final ratification.

The final recommendation statement and evidence summary are published at the same time in a peer-reviewed journal. All recommendation statements and supporting evidence reports are made available on the Task Force Web site (www.USPreventiveServicesTaskForce.org).

Please visit the Task Force Web site to learn how and when to nominate topics for consideration by the Task Force or to comment on topics in development.

Identifying High Priority Research Gaps

In the Patient Protection and Affordable Care Act of 2010, Congress has specifically charged the Task Force with identifying and reporting each year on areas where current evidence is insufficient to make a recommendation on the use of a clinical preventive service, with special attention to those areas where evidence is needed to make recommendations for specific populations and age groups. The USPSTF believes that identifying evidence gaps and highlighting them as priority areas for research will inspire public and private researchers to collaborate and target their efforts to generate new knowledge and address important health priorities. The Annual Report to Congress is available on the Task Force Web site.

Online Resources

On the Task Force Web site, people can:

- View all current USPSTF recommendations, EPC reports, and supporting materials.
- View manuals, slides, videos, and commentaries about the methods and processes the Task Force uses.
- Learn how to interpret recommendations and use them in clinical primary care practice.
- Nominate a new USPSTF member or a topic for a consideration by the Task Force.
- Provide input on specific draft materials during public comment periods.
- Sign up for the USPSTF listserv to receive USPSTF updates.

- Access the Electronic Preventive Services Selector (ePSS), a quick hands-on tool designed to help primary care clinicians and health care teams identify, prioritize, and offer the screening, counseling, and preventive medication services that are appropriate for their patients. The ePSS is available on the Web (epss.ahrq.gov) or as a mobile phone or PDA application.
- Access myhealthfinder. myhealthfinder is a tool for consumers that provides personalized recommendations for preventive services based on the U.S. Preventive Services Task Force; the *Bright Futures* Guidelines; the Centers for Disease Control and Prevention's Advisory Committee on Immunization Practices (ACIP); and the Institute of Medicine's (IOM's) Committee on Preventive Services for Women.

Steps the USPSTF Takes to Make a Recommendation



Appendix E

More Resources



AHRQ's Improving Primary Care Program

AHRQ's Improving Primary Care Program Web site (www.ahrq.gov/improvingprimarycare) produces evidence and tools to transform the primary care delivery system, resulting in better care, better experiences, and better value. The Program's Web site includes tools, resources, and materials to support health care organizations and engage the entire health care delivery system.

The Program includes two overall project areas with specific areas of focus:

- Improving Primary Care Practice
 - Capacity building
 - Care coordination
 - Clinical-community linkages
 - Health care/system redesign
 - Health information technology integration
 - Behavioral and mental health
 - Primary care practice-based research networks
 - Self-management support
- Evidence-Based Decisionmaking
 - Clinical decision support
 - Multiple chronic conditions
 - Research centers for excellence in clinical preventive services
 - U.S. Preventive Services Task Force



myhealthfinder

A consumer-friendly resource, myhealthfinder (available at www.healthfinder.gov) helps people create a customized list of relevant recommendations for preventive services based on age, sex, and pregnancy status, along with explanations of each recommendation in plain language.



Stay Healthy Brochures

Consumers can use the information in this series of brochures to learn which screening tests you need and when to get them, which medicines may prevent diseases, and daily steps to take for good health. The series includes *Men Stay Healthy at Any Age*, *Women Stay Healthy at Any Age*, *Men Stay Healthy at 50+*, and *Women Stay Healthy at 50+*, all in English and Spanish. Go to <http://www.ahrq.gov/patients-consumers/prevention/lifestyle/index.html> for the list and choose the title you are interested in.



Community Preventive Services Task Force

Established in 1996 by the U.S. Department of Health and Human Services, the Community Preventive Services Task Force (CPSTF) complements the work of the USPSTF, by addressing preventive services at the community level. The CPSTF assists agencies, organizations, and individuals at all levels (national, State, community, school, worksite, and health care system) by providing evidence-based recommendations about community prevention programs and policies that are effective in saving lives, increasing longevity, and improving Americans' quality of life. The recommendations of the CPSTF are available at www.thecommunityguide.org.



Advisory Committee on Immunization Practices

The Advisory Committee on Immunization Practices, managed and supported by the Centers for Disease Control and Prevention, is a group of medical and public health experts that develops recommendations on how to use vaccines to control diseases in the United States. The recommendations stand as public health advice that will lead to a reduction in the incidence of vaccine preventable diseases and an increase in the safe use of vaccines and related biological products.



The Substance Abuse and Mental Health Services Administration

The Substance Abuse and Mental Health Services Administration provides resources that help health care providers locate and utilize behavioral health services. The Treatment Locator (findtreatment.samhsa.gov) helps find alcohol and drug abuse treatment or mental health treatment facilities and programs around the country, and the National SBIRT Addiction Technology Transfer Center (ireta.org/ebpsbirt) helps utilize the public health model of screening and counseling to patients on risky or harmful drug and alcohol use.





Healthy People 2020

Healthy People 2020 is an initiative from the U.S. Department of Health and Human Services that challenges individuals, communities, and professionals to take specific steps to ensure good health. Healthy People provides science-based, 10-year national objectives for improving the health of all Americans. Read more at www.healthypeople.gov/2020/default.aspx.



National Guideline Clearinghouse™

A public resource for evidence-based clinical practice guidelines, NGC (guideline.gov/index.aspx) was originally created by AHRQ in partnerships with the American Medical Association and the American Association of Health Plans (now America's Health Insurance Plans). The NGC mission is to provide physicians and other health professionals, health care providers, health plans, integrated delivery systems, purchasers, and others an accessible mechanism for obtaining objective, detailed information on clinical practice guidelines and to further their dissemination, implementation, and use of this information.



Canadian Task Force on Preventive Health Care

The Task Force was established by the Public Health Agency of Canada to develop clinical practice guidelines that support primary care providers in delivering preventive health care. The mandate of the Task Force is to develop and disseminate clinical practice guidelines for primary and preventive care, based on systematic analysis of scientific evidence. Read more at www.canadiantaskforce.ca/.



Cancer Control P.L.A.N.E.T.

A service of the National Cancer Institute, the Cancer Control P.L.A.N.E.T. portal provides access to Web-based resources that can help planners, program staff, and researchers to design, implement, and evaluate evidence-based cancer control programs. Read more at cancercontrolplanet.cancer.gov/index.html.



HealthCare.gov

This Web site (www.healthcare.gov), managed by the U.S. Department of Health and Human Services, helps people take health care into their own hands. It provides information about insurance options, using insurance, the Affordable Care Act, comparing providers, and prevention and wellness—including which preventive services are covered under the Act.

Appendix F

Breast Cancer (2002 Recommendation)*

| Screening for Breast Cancer (2002 Recommendation) | | | |
|---|--|--|--|
| Title | Screening for Breast Cancer (2002 Recommendation) | | |
| Population | Women ages 40 years and older | | |
| Screening Test | Mammography, with or without clinical breast examination | Clinical breast examination alone | Breast self-examination alone |
| Recommendation | Screen every 1 to 2 years. Grade: B | No recommendation. Grade: I (Insufficient Evidence) | No recommendation. Grade: I (Insufficient Evidence) |
| Risk Assessment | <p>Women who are at increased risk for breast cancer (e.g., those with a family history of breast cancer in a mother or sister, a previous breast biopsy revealing atypical hyperplasia, or first childbirth after age 30) are more likely to benefit from regular mammography than women at lower risk.</p> <p>There is fair evidence that mammography screening every 12 to 33 months significantly reduces mortality from breast cancer. Evidence is strongest for women ages 50 to 69 years. For women ages 40 to 49 years, the evidence that screening mammography reduces mortality from breast cancer is weaker, and the absolute benefit of mammography is smaller, than it is for older women.</p> <p>Clinicians should refer patients to mammography screening centers with proper accreditation and quality assurance standards to ensure accurate imaging and radiographic interpretation. Clinicians should adopt office systems to ensure timely and adequate follow-up of abnormal results.</p> | | |
| Screening Tests | <p>The precise age at which the benefits from screening mammography justify the potential harms is a subjective judgment and should take into account patient preferences. Clinicians should inform women about the potential benefits (reduced chance of dying from breast cancer), potential harms (false-positive results, unnecessary biopsies), and limitations of the test that apply to women their age. The balance of benefits and potential harms of mammography improves with increasing age for women ages 40 to 70 years.</p> <p>Clinicians who advise women to perform breast self-examination or who perform routine clinical breast examination to screen for breast cancer should understand that there is currently insufficient evidence to determine whether these practices affect breast cancer mortality, and that they are likely to increase the incidence of clinical assessments and biopsies.</p> | | |
| Balance of Benefits and Harms | <p>The USPSTF has made recommendations on screening for genetic susceptibility for breast cancer and chemoprevention of breast cancer. These recommendations are available at http://www.uspreventiveservicestaskforce.org/.</p> | | |
| Other Relevant USPSTF Recommendations | <p>The U.S. Department of Health and Human Services, in implementing the Affordable Care Act, under the standard it sets out in revised Section 2713(a)(6) of the Public Health Service Act, utilizes the 2002 recommendation on breast cancer screening of the U.S. Preventive Services Task Force.</p> <p>For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to http://www.uspreventiveservicestaskforce.org/.</p> | | |

Index

| | |
|--|--------|
| 2-hour postload plasma (see Diabetes Mellitus, Screening) | |
| AAA (see Abdominal Aortic Aneurysm, Screening) | |
| <i>Abdominal Aortic Aneurysm, Screening</i> | 7, 93 |
| *Alcohol Misuse, Screening and Behavioral Counseling | 8 |
| Anemia, Iron Deficiency (see Iron Deficiency Anemia, Screening) | |
| Ankle Brachial Index (see Peripheral Artery Disease And Cardiovascular Risk Assessment, Screening) | |
| <i>Aspirin for the Prevention of Cardiovascular Disease, Preventive Medication</i> | 9, 93 |
| <i>Aspirin or NSAIDs for Prevention of Colorectal Cancer, Preventive Medication</i> | 10, 93 |
| <i>Aspirin Use to Prevent Preeclampsia, Preventive Medication</i> | 93 |
| Asymptomatic Bacteriuria (see Bacteriuria, Screening) | |
| <i>Autism Spectrum Disorder in Young Children, Screening</i> | 93 |
| Autorefraction (see Visual Impairment in Children Ages 1-5, Screening) | |
| Bacterial Vaginosis in Pregnancy, Screening | 11 |
| Bacteriuria, Screening | 12 |
| Basal Cell Cancer (see Skin Cancer, Counseling OR Screening) | |
| Bladder Cancer, Screening | 13 |
| Blood Lead Levels in Children and Pregnant Women, Screening | 67 |
| Blood Pressure, High (see High Blood Pressure in Adults, Screening) | |
| BMI Screening, Children and Adolescents (see Obesity in Children and Adolescents, Screening) | |
| Bone Mineral Density (see Osteoporosis, Screening) | |
| BRCA Mutation Testing (see BRCA-Related Cancer in Women, Screening) | |
| *BRCA-Related Cancer in Women, Screening | 14 |
| *Breast Cancer, Preventive Medications | 15 |
| <i>Breast Cancer, Screening</i> | 16, 94 |
| Breast Self Examination [BSE] (see Breast Cancer, Screening) | |
| Breastfeeding, Counseling | 17 |
| CA-125 Screening for Ovarian Cancer (see BRCA-Related Cancer in Women, Screening) | |

| | |
|--|--------|
| Calcium Supplements (see Vitamin D and Calcium Supplementation to Prevent Fractures, Preventive Medication) | |
| Cancer | |
| (see Aspirin or NSAIDS for Prevention of Colorectal Cancer, Preventive Medication) | |
| (see Bladder Cancer, Screening) | |
| (see BRCA-Related Cancer in Women, Screening) | |
| (see Breast Cancer, Preventive Medications) | |
| (see Breast Cancer, Screening) | |
| (see Cervical Cancer, Screening) | |
| (see Colorectal Cancer, Screening) | |
| (see Lung Cancer Screening) | |
| (see Oral Cancer, Screening) | |
| (see Ovarian Cancer, Screening) | |
| (see Prostate Cancer, Screening) | |
| (see Skin Cancer, Counseling) | |
| (see Skin Cancer, Screening) | |
| (see Testicular Cancer, Screening) | |
| <i>Carotid Artery Stenosis, Screening</i> | 18, 93 |
| Cervical Cancer, Screening | 19 |
| Chest X-Ray (see Lung Cancer, Screening) | |
| *Child Maltreatment, Interventions to Prevent | 68 |
| <i>Chlamydial Infection, Screening</i> | 20, 93 |
| Chronic Bilirubin Encephalopathy (see Hyperbilirubinemia in Infants, Screening) | |
| *Chronic Kidney Disease, Screening | 21 |
| <i>Chronic Obstructive Pulmonary Disease, Screening</i> | 22, 94 |
| Clinical Breast Examination [CBE] (see Breast Cancer, Screening) | |
| *Cognitive Impairment in Older Adults, Screening | 23 |
| Colonoscopy (see Colorectal Cancer, Screening) | |
| Colorectal Cancer, Aspirin/NSAIDS (see Aspirin or NSAIDS for Prevention of Colorectal Cancer, Preventive Medication) | |
| <i>Colorectal Cancer, Screening</i> | 24, 94 |
| Congenital Hypothyroidism, Screening | 69 |
| Coronary Heart Disease Prevention (see Aspirin for the Prevention of Cardiovascular Disease, Preventive Medication) | |

| | |
|--|--------|
| Coronary Heart Disease (Risk Assessment, Nontraditional Risk Factors), Screening | 25 |
| *Coronary Heart Disease, Screening With Electrocardiography | 26 |
| COPD (see Chronic Obstructive Pulmonary Disease, Screening) | |
| CT (Computerized Tomography) for Lung Cancer (See Lung Cancer, Screening) | |
| Cover-Uncover Test (see Visual Impairment in Children Ages 1-5, Screening) | |
| Dementia (see Cognitive Impairment in Older Adults, Screening) | |
| <i>Dental Caries Prevention in Children from Birth to Age 5 Years, Screening and Preventive Medication</i> | 93 |
| <i>Depression in Adults, Screening</i> | 27, 94 |
| Depression or Depressive Disorders in Children and Adolescents (see Major Depressive Disorder in Children and Adolescents, Screening) | |
| <i>Depression in Pregnant and Postpartum Women, Screening</i> | 94 |
| Developmental Dysplasia of the Hip, Screening | 70 |
| <i>Diabetes Mellitus, Screening</i> | 28, 93 |
| Drug Use, Illicit (see Illicit Drug Use, Screening) | |
| Drug Abuse (see Illicit Drug Use, Screening) | |
| Dysplasia, Hip (see Developmental Dysplasia of the Hip, Screening) | |
| Elevated Blood Lead Levels (see Blood Lead Levels in Children and Pregnant Women, Screening) | |
| Estrogen Therapy (see Menopausal Hormone Therapy for the Primary Prevention of Chronic Conditions, Preventive Medication) | |
| *Falls in Older Adults, Counseling, Preventive Medication, and Other Interventions | 29 |
| Fasting Plasma Glucose (see Diabetes Mellitus, Screening) | |
| Fecal Occult Blood Testing [FOBT] (see Colorectal Cancer, Screening) | |
| Folic Acid Supplementation to Prevent Neural Tube Defects, Preventive Medication | 30 |
| Framingham-based Adult Treatment Panel III (see Lipid Disorders in Adults, Screening or Healthful Diet and Physical Activity, Counseling) | |
| Genetic Risk Assessment and BRCA Mutation Testing (see BRCA-Related Cancer in Women, Screening) | |
| Genital Herpes, Screening | 31 |
| *Gestational Diabetes Mellitus, Screening | 32 |
| *Glaucoma, Screening | 33 |

| | |
|--|--------|
| Glucose Challenge, Glucose Tolerance (see Gestational Diabetes Mellitus, Screening) | |
| Gonococcal Ophthalmia Neonatorum, Preventive Medication | 71 |
| Gonorrhea, Screening | 34, 93 |
| *Healthful Diet and Physical Activity, Counseling | 35 |
| <i>Healthy Diet and Physical Activity for Cardiovascular Disease Prevention in Adults at Risk, Counseling</i> | 93 |
| Hearing Loss in Newborns, Screening | 72 |
| *Hearing Loss in Older Adults, Screening | 36 |
| Heart Disease (see Coronary Heart Disease, Risk Assessment, Nontraditional Risk Factors) | |
| Hemochromatosis, Screening | 37 |
| Hemoglobin A1C (see Diabetes Mellitus, Screening) | |
| <i>Hepatitis B Virus Infection, Screening</i> | 94 |
| Hepatitis B Virus Infection in Pregnant Women, Screening | 38 |
| *Hepatitis C Virus Infection in Adults, Screening | 39 |
| Hereditary Hemochromatosis (see Hemochromatosis, Screening) | |
| Herpes Simplex Virus (see Genital Herpes, Screening) | |
| High Blood Pressure in Adults, Screening | 40, 93 |
| *High Blood Pressure in Children, Screening | 73 |
| High Cholesterol (see Lipid Disorders in Adults OR in Children, Screening) | |
| Hip Dysplasia (see Developmental Dysplasia of the Hip, Screening) | |
| *HIV Infection, Screening | 41 |
| Hirschberg Light Reflex Test (see Visual Impairment in Children Ages 1-5, Screening) | |
| Hormone Replacement Therapy (see Menopausal Hormone Therapy for the Primary Prevention of Chronic Conditions, Preventive Medication) | |
| HPV Testing (see Cervical Cancer, Screening) | |
| HT or HRT (see Menopausal Hormone Therapy for the Primary Prevention of Chronic Conditions, Preventive Medication) | |
| Human Immunodeficiency Virus (see HIV Infection, Screening) | |
| Hyperbilirubinemia in Infants, Screening | 74 |
| Hyperlipidemia (see Lipid Disorders in Adults OR in Children, Screening) | |
| Hypertension (see High Blood Pressure in Adults OR in Children, Screening) | |
| Hypothyroidism, Congenital (see Congenital Hypothyroidism, Screening) | |
| Illicit Drug Use, Screening | 42 |

| | |
|---|------------|
| *Illicit and Prescription Drug Use in Children and Adolescents, Counseling | 75 |
| Impaired Visual Acuity in Older Adults, Screening | 43, 94 |
| *Intimate Partner Violence and Elderly Abuse, Screening | 44 |
| <i>Iron Deficiency Anemia, Screening</i> | 76, 77, 93 |
| Iron Supplementation (see Iron Deficiency Anemia, Screening) | |
| Lead Levels in Blood, Elevated (see Blood Lead Levels in Children and Pregnant Women, Screening) | |
| <i>Lipid Disorders in Adults, Screening</i> | 45, 94 |
| <i>Lipid Disorders in Children, Screening</i> | 78, 94 |
| Low Dose Computerized Tomography (see Lung Cancer, Screening) | |
| *Lung Cancer, Screening | 46 |
| <i>Major Depressive Disorder in Children and Adolescents, Screening</i> | 79, 93 |
| Major Depressive Disorder in Adults (see Depression in Adults, Screening) | |
| Mammography (see Breast Cancer, Screening) | |
| Melanoma (see Skin Cancer, Counseling OR Screening) | |
| *Menopausal Hormone Therapy for the Primary Prevention of Chronic Conditions, Preventive Medication | 47 |
| Motor Vehicle Occupant Restraints, Counseling | 48 |
| Multivitamins (see Vitamin, Mineral, and Multivitamin Supplements for the Primary Prevention of Cardiovascular Disease and Cancer, Preventive Medication) | |
| Neural Tube Defects (see Folic Acid Supplementation to Prevent Neural Tube Defects, Preventive Medication) | |
| Newborn Hearing Screening (see Hearing Loss in Newborns, Screening) | |
| Nonmedical Prescription Drug Use in Children and Adolescents (see Illicit and Prescription Drug Use in Children and Adolescents, Counseling) | |
| Non-Steroidal Anti-Inflammatories [NSAIDs] (see Aspirin or NSAIDs for Prevention of Colorectal Cancer, Preventive Medication) | |
| *Obesity in Adults, Screening | 49 |
| Obesity in Children and Adolescents, Screening | 80 |
| *Oral Cancer, Screening | 50 |
| Osteoporosis, Screening | 51 |
| *Ovarian Cancer, Screening | 52 |
| Ovarian Cancer, Genetic Risk (see BRCA-Related Cancer in Women, Screening) | |
| Overweight (see Obesity in Children and Adolescents, Screening) | |
| Pap Smear (see Cervical Cancer, Screening) | |

| | |
|---|--------|
| *Peripheral Artery Disease and Cardiovascular Risk Assessment, Screening | 53 |
| Phenylketonuria (PKU), Screening | 81 |
| Photoscreening (see Visual Impairment in Children Ages 1-5, Screening) | |
| PKU (see Phenylketonuria, Screening) | |
| Postmenopausal Hormone Therapy (see Menopausal Hormone Therapy for the Primary Prevention of Chronic Conditions, Preventive Medication) | |
| Prescription Drug Abuse (see Illicit and Prescription Drug Use in Children and Adolescents) | |
| Progestin Therapy (see Menopausal Hormone Therapy for the Primary Prevention of Chronic Conditions, Preventive Medication) | |
| *Prostate Cancer, Screening | 54 |
| PSA Screening for Prostate Cancer (see Prostate Cancer, Screening) | |
| <i>Sexually Transmitted Infections, Counseling</i> | 55, 93 |
| Sickle Cell Disease in Newborns, Screening | 82 |
| Sigmoidoscopy (see Colorectal Cancer, Screening) | |
| *Skin Cancer, Counseling | 56 |
| <i>Skin Cancer, Screening</i> | 57, 94 |
| <i>Speech and Language Delay, Screening</i> | 83, 94 |
| Squamous Cell Cancer (see Skin Cancer, Counseling OR Screening) | |
| Smoking Cessation (see Tobacco Use in Adults, Counseling and Interventions OR Tobacco Use in Children and Adolescents, Primary Care Interventions) | |
| Spirometry Screening for COPD (see Chronic Obstructive Pulmonary Disease, Screening) | |
| Sputum Cytology (see Lung Cancer, Screening) | |
| Stereoacuity Test (see Visual Impairment in Children Ages 1-5, Screening) | |
| STI or STD | |
| (see Sexually Transmitted Infections, Counseling) | |
| (see Chlamydial Infection, Screening) | |
| (see Gonorrhea, Screening) | |
| (see Genital Herpes, Screening) | |
| (see HIV Infection, Screening) | |
| (see Syphilis Infection (Pregnant Women), Screening) | |
| <i>Suicide Risk, Screening</i> | 58, 94 |
| <i>Syphilis Infection, Screening</i> | 94 |

| | |
|---|--------|
| Syphilis Infection (Pregnant Women), Screening | 59 |
| T4 (see Congenital Hypothyroidism, Screening) | |
| Testicular Cancer, Screening | 60 |
| <i>Thyroid Disease Screening</i> | 93 |
| *Tobacco Use in Children and Adolescents, Primary Care Interventions | 84 |
| <i>Tobacco Use in Adults, Counseling and Interventions</i> | 61, 94 |
| TSH (see Congenital Hypothyroidism, Screening) | |
| Ultrasonography (see Abdominal Aortic Aneurysm, Screening) | |
| Urinalysis (see Bladder Cancer, Screening) | |
| Urine Biomarkers (see Bladder Cancer, Screening) | |
| Urine Culture (see Bacteriurias, Screening) | |
| Urine Cytology (see Bladder Cancer, Screening) | |
| Vaginosis, Bacterial (see Bacterial Vaginosis in Pregnancy, Screening) | |
| Visual Acuity Test (see Visual Impairment in Children Ages 1-5, Screening) | |
| Visual Impairment in Children Ages 1-5, Screening | 85 |
| *Vitamin D and Calcium Supplementation to Prevent Fractures, Preventive Medication | 62 |
| <i>Vitamin D Deficiency, Screening</i> | 94 |
| Vitamin D3 Supplement (see Vitamin D and Calcium Supplementation to Prevent Fractures, Preventive Medication or Falls in Older Adults) | |
| *Vitamin, Mineral, and Multivitamin Supplements for the Primary Prevention of Cardiovascular Disease and Cancer, Preventive Medication | 63 |

* indicates new or updated recommendations released March 2012 to March 2014.

Bold text indicates topic of recommendation.

Italic text indicates topic in progress (new or update).

AHRQ's Electronic Preventive Services Selector (ePSS)

**Bringing the prevention information clinicians need—
recommendations, clinical considerations, and selected practice
tools—to the point of care.**

The ePSS helps you identify and select screening, counseling, and preventive medication services based on specific patient characteristics.



Available at

<http://epss.ahrq.gov>

on the Web and for mobile devices, including Android, BlackBerry,
iPhone, Palm, and iPad.

The U.S. Preventive Services Task Force is always interested in making our resources and tools more useful to those implementing the recommendations in primary care, as well as educating health professions students and patients. Please send any comments to coordinator@USPSTF.net.



U.S. Department of Health and Human Services
Agency for Healthcare Research and Quality

www.ahrq.gov

AHRQ Pub. No. 14-05158

May 2014

ISBN 978-1-58763-439-0